



**Canadian  
Intellectual Property  
Office**

An Agency of  
Industry Canada

**Office de la Propriété  
Intellectuelle  
du Canada**

Un organisme  
d'Industrie Canada

(11) **CA 2 447 750**

(13) **A1**

(40) 12.12.2002

(43) 12.12.2002

(12)

(21) **2 447 750**

(22) **27.05.2002**

(51) Int. Cl. 7:

**C12Q 1/68**, C12N 15/31,  
G01N 33/53, G01N 33/566,  
G01N 33/569

(85) **18.11.2003**

(86) **PCT/JP02/005107**

(87) **WO02/099133**

(30) **2001-165929 JP 31.05.2001**

(71)

**FUSO PHARMACEUTICAL INDUSTRIES, LTD.,  
7-10, Dosho-machi 1-chome  
Chuo-ku, Osaka-shi  
541-0045, OSAKA, XX (JP).  
OHNO, TSUNEYA,  
5-15 Fukazawa 2-chome**

**Setagaya-ku  
158-0081, TOKYO, XX (JP).**

(72)

**MATSUHIRA, AKIO (JP).  
OHNO, TSUNEYA (JP).**

(74)

**KIRBY EADES GALE BAKER**

(54) **PROCEDE AMELIORE DE DETECTION ET D'IDENTIFICATION DE MICRO-ORGANISMES CAUSES D'INFECTIONS**

(54) **IMPROVED METHOD FOR DETECTING AND IDENTIFYING MICROORGANISM CAUSATIVE OF INFECTION**

(57)

A microorganism causative of an infection is quickly and highly sensitively detected and/or identified by obtaining phagocytes from a clinical specimen containing phagocytes originating in a living body, immobilizing the obtained phagocytes, treating the phagocytes so as to enhance the permeability of the phagocyte membrane and expose DNA of the microorganism causative of the infection which is anticipated as being contained in the phagocytes, and then using a detection DNA probe hybridizable with the DNA under stringent conditions.



Office de la Propriété  
Intellectuelle  
du Canada

Un organisme  
d'Industrie Canada

Canadian  
Intellectual Property  
Office

An agency of  
Industry Canada

CA 2447750 A1 2002/12/12

(21) **2 447 750**

(12) **DEMANDE DE BREVET CANADIEN  
CANADIAN PATENT APPLICATION**

(13) **A1**

(86) Date de dépôt PCT/PCT Filing Date: 2002/05/27  
(87) Date publication PCT/PCT Publication Date: 2002/12/12  
(85) Entrée phase nationale/National Entry: 2003/11/18  
(86) N° demande PCT/PCT Application No.: JP 2002/005107  
(87) N° publication PCT/PCT Publication No.: 2002/099133  
(30) Priorité/Priority: 2001/05/31 (2001-165929) JP

(51) Cl.Int.<sup>7</sup>/Int.Cl.<sup>7</sup> C12Q 1/68, C12N 15/31, G01N 33/53,  
G01N 33/569, G01N 33/566

(71) Demandeurs/Applicants:  
FUSO PHARMACEUTICAL INDUSTRIES, LTD., JP;  
OHNO, TSUNEYA, JP

(72) Inventeurs/Inventors:  
MATSUHISA, AKIO, JP;  
OHNO, TSUNEYA, JP

(74) Agent: KIRBY EADES GALE BAKER

(54) Titre : PROCÉDE AMÉLIORÉ DE DÉTECTION ET D'IDENTIFICATION DE MICRO-ORGANISMES CAUSES  
D'INFECTIONS

(54) Title: IMPROVED METHOD FOR DETECTING AND IDENTIFYING MICROORGANISM CAUSATIVE OF  
INFECTION

(57) Abrégé/Abstract:

A microorganism causative of an infection is quickly and highly sensitively detected and/or identified by obtaining phagocytes from a clinical specimen containing phagocytes originating in a living body, immobilizing the obtained phagocytes, treating the phagocytes so as to enhance the permeability of the phagocyte membrane and expose DNA of the microorganism causative of the infection which is anticipated as being contained in the phagocytes, and then using a detection DNA probe hybridizable with the DNA under stringent conditions.

Canada

<http://opic.gc.ca> • Ottawa-Hull K1A 0C9 • <http://cipo.gc.ca>

OPIC • CIPQ 191

OPIC



CIPQ

**ABSTRACT**

Causative microorganisms of infectious diseases are detected and/or identified rapidly and high-sensitively by taking phagocytes from the clinical specimens containing active phagocytes, immobilizing the phagocytes so taken, treating the phagocytes to improve cell membrane permeabilities thereof, further treating the phagocytes to bare DNA in the causative microorganisms which might be existed in the phagocytes, and detecting the causative microorganisms with DNA probes which can hybridize with such DNA under stringent conditions.

## **SPECIFICATION**

### **IMPROVED METHOD FOR DETECTING AND IDENTIFYING CAUSATIVE MICROORGANISMS OF INFECTIOUS DISEASES**

#### **TECHNICAL FIELD**

The present invention relates to an improved method for detecting and identifying causative microorganisms of infectious diseases. The present invention also relates to a kit for detecting and/or identifying causative microorganisms of infectious diseases, a method for monitoring genes from exogenous microorganisms in clinical specimens, and a method for determining causative microorganisms of sepsis and those of bacteriemia.

#### **BACKGROUND ARTS**

Although the hemoculture methodologies have popularly been used conventionally as a mean to verify bacteria in the blood, since this methodology needs about from 3 to 14 days to culture and isolate the subjected bacteria and detection rates thereby are as low as about 10%, it was not well contributed in the diagnosis for treating serious diseases like sepsis.

The present inventors had invented, to solve such problems, a method for detecting and identifying exogenous-microorganisms digested with phagocytes comprising a step of detecting genes from such exogenous-microorganisms in the phagocytes by *in situ* hybridization employing a probe which can specifically hybridize with the genes (Japanese Patent Publication No. 7-40).

The method of Japanese Patent Publication No. 7-40 have been in the limelight in the field of infectious diseases because, in comparison with the conventional hemoculture methodology, the method allowed about four times rapidly detection of the subjected bacteria in bloods from patients who are under the suspicion about sepsis, and detection results were appeared within 24 hours.



Objects of the present inventions is an improvement of detection effects and of detection sensitivity to be offered by the method according to Japanese Patent Publication No. 7-40 for detecting and/or identifying causative microorganisms of infectious diseases by taking phagocytes from the clinical specimens containing active phagocytes, immobilizing the phagocytes so taken, treating the phagocytes to improve cell membrane permeabilities thereof, further treating the phagocytes to bare DNA in the causative microorganisms which might be existed in the phagocytes, *in situ* hybridizing DNA so bared with detective DNA probe(s) which can hybridize with such bared DNA under stringent conditions, and detecting and/or identifying the causative microorganisms based on signals so detected.

#### DISCLOSURE OF INVENTION

The present invention has been completed in view of the problems aforementioned and the merits thereof are as follows.

A method for detecting and/or identifying causative microorganisms of infectious diseases by taking phagocytes from the clinical specimens containing active phagocytes, immobilizing the phagocytes so taken, treating the phagocytes to improve cell membrane permeabilities thereof, further treating the phagocytes to bare DNA in the causative microorganisms which might be existed in the phagocytes, *in situ* hybridizing DNA so bared with detective DNA probe(s) which can hybridize with such bared DNA under stringent conditions, and detecting and/or identifying the causative microorganisms based on signals so detected, the method comprises at least one condition(s) to be selected from the following conditions (1)-(8) of;

(1) Cell density ( $X$  cells/ml) of the phagocytes to be immobilized is  $5 \times 10^6$  cells/ml  $< X$  cells/ml  $< 1 \times 10^8$  cells/ml,

(2) Lysostaphin is applied into the step to bare DNA in the titer of from 1 Unit/ml to 1,000 Units/ml,

(3) Lysozyme is applied into the step to bare DNA in the titer of from 1,000 Units/ml to 1,000,000 Units/ml,

(4) N-acetylmuramidase is applied into the step to bare DNA in the titer of from 10 Units/ml to 10,000 Units/ml,

(5) Zymolyase is applied into the step to bare DNA in the titer of from 50 Units/ml to 500 Units/ml,

(6) Surfactant is applied into the step of *in situ* hybridization,

(7) Such DNA probe(s) is/are one or more DNA probe(s) to be determined with their chain length of from 350 bases to 600 bases, and

(8) Concentration of such DNA probe(s) is from 0.1 ng/ $\mu$ l to 2.2 ng/ $\mu$ l.

The step to bare DNA employs preferably one or more enzyme(s) selected from Lysostaphin in the titer of from about 10 to about 100 Units/ml, Lysozyme in the titer of from about 10,000 to about 100,000 Units/ml, N-acetylmuramidase in the titer of from about 100 to about 1,000 Units/ml and Zymolyase in the titer of from about 100 to about 500 Units/ml.

The step to bare DNA preferably employs enzyme(s), and the enzyme(s) is/are subjected to a reaction to be performed under the temperature of from about 26°C to about 59°C for from about 15 to about 120 minutes.

The step to bare DNA further employs preferably substance(s), in particular, phenylmethanesulfonyl fluoride to keep a form of the phagocytes under the concentration of preferably from about 10 $\mu$ mol/l to about 10mmol/l.

As the substance to keep a form of the phagocytes, substance dissolved into dimethylsulfoxide is preferable. The substance dissolved in dimethylsulfoxide is employed as that to keep a form of the phagocytes, then, concentration of the dimethylsulfoxide in a solution to be used in the step to bare DNA is adjusted to less than 5%.

The *in situ* hybridization step is performed by hybridizing DNA with DNA probe(s) under the presence of surfactant(s), in particular, the anionic surfactant, preferably, sodium dodecyl sulfate (SDS).

Hybridization reaction in the *in situ* hybridization step is performed under the temperature of from about 20°C to about 50°C and time of from about 30 to about 900 minutes.

The method further comprises, prior to the immobilization step, a step to mount the phagocytes so taken onto the solid support and a slide coated with 3-aminopropyl triethoxysilane is employed as such solid support.

Pigment(s) is/are also employed at the signal detection to distinguish contrast between signals and cells. Preferably, blood is employed as the clinical specimen.

The present invention further provides a kit for detecting and/or identifying causative microorganisms of infectious diseases by taking phagocytes from the clinical specimens containing active phagocytes, immobilizing the phagocytes so taken, treating the phagocytes to improve cell membrane permeabilities thereof, further treating the phagocytes to bare DNA in the causative microorganisms which might be existed in the phagocytes, *in situ* hybridizing DNA so bared with detective DNA probe(s) which can hybridize with such bared DNA under stringent conditions, and detecting and/or identifying the causative microorganisms based on signals so detected, the kit comprises the following elements (1)-(2) of;

(1) at least one enzyme(s) to be employed in the step to bare DNA which is/are selected from Lysostaphin, Lysozyme, N-acetylmuramidase and Zymolyase; and

(2) at least one detective DNA probe(s).

Then, the present invention further provides a method for monitoring a gene of exogenous microorganisms digested with the phagocytes in the clinical specimens containing active phagocytes comprising the step of detecting the gene with *in situ* hybridization method employed in the foregoing method, wherein the gene of exogenous microorganisms in the clinical specimens is monitored.

The present invention further provides a method for diagnosing sepsis or bacteriemia comprising the step of identifying a gene of candidate causative microorganisms with *in situ* hybridization method employed in the foregoing method, wherein the causative microorganisms for sepsis or bacteriemia are determined based on the identification results.

#### BRIEF DESCRIPTION OF DRAWINGS

Figure 1 is a view of illustrating results of *in situ* hybridization performed under (a) the absence of surfactant (SDS) and (b) the presence of surfactant (SDS).

Figure 2 is a view of illustrating a manner of leukocytes immobilized with the various cell density.

Figure 3 is a view of illustrating a time-coursely change on lytic enzyme activity against (a) *Staphylococcus aureus* and *Staphylococcus epidermidis*, (b) *Pseudomonas aeruginosa* and *Escherichia coli*, and (c) *Enterococcus faecalis*.

Figure 4 is a view of illustrating concentration-dependency effects by addition of DMSO against lytic activity to be offered by (a) 300 Units/ml of N-acetylmuramidase, (b) 10,000 Units/ml of Lysozyme, and (c) 50 Units/ml of Lysostaphin.

Figure 5 is a view of illustrating results on addition of (a) protease 0.2 Units/ml only, (b) PMSF 1  $\mu$ mol/ml, (c) PMSF 10  $\mu$ mol/ml, (d) PMSF 0.1mmol/ml, and (e) PMSF 1mmol/ml to study effects of PMSF to be used to suppress the function of protease

which changes a morphological form of leukocyte.

Figure 6 is a view of illustrating that, in the phagocytosis samples prepared according to the present invention, phagocytes digested bacteria and morphological forms of such digested bacteria were changed.

Figure 7 is a view of illustrating effects of enzyme-treatment in the phagocytosis samples and manner of the phagocytosis samples containing (a) *Staphylococcus aureus* prior to the treatment, (b) *Enterococcus faecalis* prior to the treatment, (c) *Staphylococcus aureus* of (a) with the treatment, and (d) *Enterococcus faecalis* of (b) with the treatment.

Figure 8 is a shematic view of the slide on which the phagocytosis samples were smeared to study optimum probe concentration under the *in situ* hybridization.

Figure 9 is a shematic view of the slide on which the phagocytosis samples were smeared to study optimum temperature under the *in situ* hybridization.

Figure 10 is a view of illustrating the signals appeared in the results of Southern Blot (the upper row) and Electrophoresis (the lower row) together with the length of the labelled detective probes prepared by putting digoxigenin-labels on (a) SA probes and (b) PA probes.

Figure 11 is a view of illustrating the signals detected by *in situ* hybridization of the digested *Escherichia coli* with detective probes of (a) EC-24, (b) EC-34, (c) EC-39 and (d) the mixed probes (MIX) of the foregoing probes (a)-(c).

BEST MODE FOR CARRYING OUT THE INVENTION

Any clinical specimen containing active phagocytes is applicable as a specimen to be employed in the present embodiment and may includes fluidus like blood, histofluid, lymph, neurolymph, pus, pituita, nasal mucus or sputum. Then, active phagocytes are contained in urea, ascites or dialysate in the disorder like diabetes, nephropathia or hepatopathy and are also remained in the lotion used to wash rhinonem, bronchus, skin, various organs or bones, they can therefore also be employed as specimen of the present invention.

In addition thereto, tissues taken from skin, lung, kidney, mucosa or the like are also employed as the clinical specimen in the present invention. This is because that macrophage, which is one of phagocytes, has various forms including monocytes, alveolar macrophage, celiac macrophage, fixed macrophage, free macrophage, Hanseemann's macrophage, inflammatory macrophage, liver Kupffer cells and brain microglia cells, tissues containing those can therefore also be employed, besides blood, as the clinical specimen of the present invention. For example, causative microorganisms of nephritis can be detected and identified by collecting kidney tissues through biopsy from patients who are under the suspicion about nephritis, taking phagocytes in the tissues through digestion thereof with enzymes like trypsin to exfoliate cells, and utilizing the phagocytes so taken.

The term `phagocytes` used herein is directed to any cell which can incorporate into itself foreign objects like exogenous microorganisms and may includes, for example, macrophage, monocytes, neutrophil and eosinophil. Phagocytes line like U937 Cell, HL60 Cell or the like is also available. Exogenous microorganisms which may cause infectious diseases are microorganisms to be digested with phagocytes and may includes, for example, bacteria, mycete, virus, protozoon, parasite or the like. Bacteria may include, for example, *Staphylococcus*, *Pseudomonas*, *Enterococcus*, *Colibacillus*, *Streptococcus*,

*Pneumococcus*, *Tubercle bacillus*, *Helicobacter pylori*, *Listeria*, *Yersinia*, *Brucella* or the like. Mycete may include, for example, *Candida*, *Aspergillus*, *Actinomyces*, *Coccidioides*, *Blastomyces* or the like. Virus may include Influenza virus, Poliovirus, Herpes virus, Hepatitis virus and AIDS virus. Protozoon may include, for example, *Karyamoebina falcata*, *Trichomonas vaginalis*, Malaria, *Toxoplasma* or the like. Parasite may include, for example, *Trypanosoma* or the like. In particular, the causative microorganisms of sepsis or bacteriemia may include, for example, Gram-Positive Bacteria of *Staphylococcus* genus (*Staphylococcus aureus*, *Staphylococcus epidermidis*) and *Enterococcus* genus (*Enterococcus faecalis*, *Enterococcus faecium*, *Streptococcus pneumoniae*, *Streptococcus pyogenes*, *Streptococcus agalactiae*), Gram-Negative Bacteria like *Colibacillus*-related *Enterobacteriaceae* Family of *Escherichia coli*, *Enterobacter cloacae*, *Klebsiella pneumoniae* (*Klebsiella oxytoca*, *Serratia marcescens*, *Proteus vulgaris*, *Citrobacter freundii*), aerophilic rod of *Pseudomonas* genus (*Pseudomonas aeruginosa*), anaerobe of *Clostridium* genus (*Clostridium perfringens*), *Bacteroides* genus (*Bacteroides fragilis*) or the like. Also, *Acinetobacter calcoaceticus*, *Aeromonas hydrophila*, *Flavobacterium meningosepticum*, *Bacillus cereus* can rarely be fallen within the causative microorganisms.

Phagocytes (Leukocytes) fractions can be taken from the clinical specimen according to the conventional method. For example, about 5ml (or 10ml with a few leukocyte) of heparinized venous bloods were obtained and the bloods were mixed with the blood components separative reagent (adjusted with sterilized-purified water to be 25ml as their final volume containing 225mg of Sodium Chloride and 1.5 g of Dextran (Molecular Weight: 200,000-300,000)) in the ratio of 4:1. Leukocyte fractions (the upper layer) were then obtained by leaving them at from about 10°C to about 40°C for from about 15 minutes to about 120 minutes, preferably, about 37°C for about 30 minutes. Leukocytes were appeared by centrifuging the leukocyte fractions so taken in from about 100×g to about 500

× g, at from 0°C to about 20°C for about from 3 minutes to about 60 minutes, preferably, in from about 140× g to about 180× g, at about 4°C for about 10 minutes. Hemolysis is preferable if erythroblasts are entered at this step. Pelletized leukocytes so obtained were, for example, suspended with 1 ml of sterilized-purified water and the suspension were immediately put into an isotonic state by adding thereto excessive amounts of PBS (prepared by diluting twenty-fold with sterilized-purified water the raw solution (PBS Raw Solution; hereinafter simply referred to as `PBS Raw Solution`) which have been adjusted with sterilized-purified water to be 120ml as their final volume containing 18.24 g of Sodium Chloride, 6.012 g of Sodium Monohydrogen Phosphate Didecahydrate and 1.123 g of Sodium Dihydrogen Phosphate Dihydrate), and such suspension were re-centrifuged in from about 140× g to about 180× g, at 4 °C and for about 10 minutes. Otherwise, such digested phagocytes can be adhered to the slides to be noted later through their native adhesionability without any centrifugation aforementioned.

Methodology to fix the leukocytes may include, for example, Carnoy's fixation. In particular, leukocytes are mounted onto a support (a supportive medium) which can hold the leukocytes and are immersed in Carnoy's fixative solution (prepared by mixing Ethanol, Chloroform and Acetic Acid in the volume ratio of 6:3:1) for 20 minutes and were immersed in from about 50% to about 90%, preferably, about 75% Ethanol solution for five minutes. Finally, they were completely air-dried.

Insoluble materials are preferable for the support and may include, for example, glass, metal, synthesized resin (e.g., polystyrene, polyethylene, polypropylene, polyvinyl chloride, polyester, polyacrylic ester, nylon, polyacetal and fluorine resin) and polysaccharide (e.g., cellulose and agarose).

A form of the insoluble support can be changed optionally and may include, for example, that of plate, tray, ball, fiber, rod, disk, container, cell or tube.



Particularly preferred support in the present embodiment is to employ slides. Such slide may include, for example, the slide (PRODUCTS ID. S311BL) of JAPAN AR BROWN CO., LTD. This slide (PRODUCTS ID. S311BL) has 14 circular wells of 5mm diameter. In order to improve an adhesionability of the subjected cells at the actual use, APS coated slides to be made by coating 3-aminopropyltriethoxysilane (APS, SIGMA) onto the slides are recommended. The other slides to be made by coating poly-L-lysine or gelatin are also available.

APS coated slides are prepared by putting the slides (PRODUCTS ID. S311BL) onto a holder, immersing them into the diluted neutral detergent for 30 minutes or more, removing well the detergent with tap water, then washing the slides with purified water, and drying well the same under the higher temperature (100°C or more) followed by leaving it under the room temperature. Thereafter, these slides are immersed into acetone containing 2% APS for one minute and are immediately rinsed gently with acetone then with sterilized-purified water. These slides were then air-dried. These slides are re-immersed into acetone containing from about 1% to about 10% APS for one minute, are immediately rinsed gently with acetone then sterilized-purified water, and are air-dried. Finally, these slides are dried under the temperature of from about 20°C to about 60°C, preferably of 42°C to realize the APS coated slides.

Leukocytes are mounted onto the APS coated slides preferably by smearing the leukocytes to realize the extended mono-layer thereof and drying the same. Cell population ( $X$  cells/ml) of the immobilized phagocytes should be adjusted to that of about  $5 \times 10^6$  cells/ml  $< X$  cells/ml  $<$  about  $1 \times 10^8$  cells/ml, preferably, about  $1 \times 10^7$  cells/ml  $\leq X$  cells/ml  $\leq$  about  $5 \times 10^7$  cells/ml.

Then, according to change on the phagocytes population per 1ml, leukocyte population to be immobilized in the single well of the APS coated slide ( $Y$  cells/well (diameter: 5mm)) are

adjusted that of about  $2.5 \times 10^4$  cells/well  $< y$  cells/well  $<$  about  $5 \times 10^5$  cells/well, preferably, about  $5 \times 10^4$  cells/well  $\leq y$  cells/well  $\leq$  about  $2.5 \times 10^5$  cells/well. In particular, pelletized leukocytes are prepared by centrifuging leukocyte fractions in from about  $140 \times g$  to about  $180 \times g$ , at  $4^\circ C$  and for 10 minutes, adding small amount of PBS to the pelletized leukocytes, suspending the same, and counting the population with a hemacytometer. Leukocytes were duly mounted on the APS coated slides by smearing  $5 \mu l$  of the leukocyte suspension into each well of the slides adjusted with PBS to be cell population of from about  $5 \times 10^4$  cells/well to about  $2.5 \times 10^5$  cells/well, then extending mono-layer of the leukocytes, and completely air-drying the same.

In order to accelerate permeability of the phagocyte membrane, they were immersed in PBS for from about 3 minutes to about 30 minutes, then in the solution prepared by diluting from about 2-fold to about 50-fold a pretreatment reagent (prepared by mixing 1.25 g of Saponin, 1.25 ml t-octylphenoxy-polyethoxyethanol (specific gravity of 1.068-1.075 ( $20/4^\circ C$ ), pH (5w/v%) 5.5-7.5) and 25ml PBS Raw Solution, and adjusting with sterilized- purified water to be 50ml as their final volume), and were applied to a centrifuge for from about 3 minutes to about 30 minutes.

In order to bare DNA in the causative microorganisms, an enzyme solution was prepared by adding, per single slide, 1 ml of a reagent solvent (prepared by diluting about 100-fold, with PBS, Dimethyl Sulfoxide (DMSO) containing 0.1 mol/l Phenylmethyl Sulfonylfluoride (PMSF)) to an enzyme reagent (N-acetyl-muramidase, Lysozyme and/or Lysostaphin), then 1 ml of which were dropped under the temperature of from about  $20^\circ C$  to about  $60^\circ C$ , preferably, from about  $37^\circ C$  to about  $42^\circ C$  in the wet chamber onto the area where the leukocytes were smeared and were left for from about 10 minutes to about 60 minutes. Then, they were immersed in PBS containing 0.2 mol/l Hydrochloric Acid (prepared by adding hydrochloric acid to PBS Raw Solution, diluting 20-fold

the same with sterilized-purified water and adjusting final concentration of Hydrochloric Acid to 0.2 mol/l), and were applied to a centrifuge for 3-30 minutes to accelerate thier permeability. 5% or more of DMSO concentration may lower activities of Lysozyme and Lysostaphin, DMSO concentration of less than 5% is therefore preferable. Besides PMSF, the known protease inhibitors like tosyllysinechloromethylketone (TLCK) and a combination thereof are also applicable to keep the form of phagocytes. Solvents like DMSO can be changed optionally to employ such known protease inhibitors.

With regard to the preferable titer range on each enzyme employed in the enzyme reagent, although Lysostaphin offer the substantial effect at the titer of 1 Unit/ml in the lysis of *Staphylococcus aureus*, that of 10 Units/ml or more was necessary in the lysis of *Staphylococcus epidermidis*. Optimum titer on Lysostaphin should therefore be adjusted to from about 1 Unit/ml to about 1,000 Units/ml, preferably, from about 10 Units/ml to about 100 Units/ml. When the titer of Lysozyme was about 10,000 Units/ml, there was no lysis on *Enterococcus faecalis* with N-acetylmuramidase of about 10 Units/ml or less titer. Also, when the titer of N-acetylmuramidase was about 100 Units/ml, there was no lysis with Lysozyme of about 1,000 Units/ml or less titer. Accordingly, optimum titer on N-acetylmuramidase should be adjusted to from about 10 Units/ml to about 10,000 Units/ml, preferably, from about 100 Units/ml to about 1,000 Units/ml, while that on Lysozyme acetylmuramidase should be adjusted to from about 1,000 Units/ml to about 1,000,000 Units/ml, preferably, from about 10,000 Units/ml to about 100,000 Units/ml. When the causative microorganisms are mycete like *Candida albicans*, titer on Zymolase should be adjusted to from about 50 Units/ml to about 500 Units/ml, preferably, from about 100 Units/ml to about 500 Units/ml. In particular, PMSF or the known protease inhibitors are useful in combination with Zymolase.

Enzyme(s) could be selected based on the components difference between Gram-Positive Bacteria and Gram-Negative Bacteria, namely, the difference of peptidoglycan or of lipopolysaccharides. Two kinds or more of enzymes are particularly preferable to effectively lyse both Gram-Positive Bacteria and Gram-Negative Bacteria. It was demonstrated in the present invention that the lysis activities offered by the mixed enzymes of Lysozyme, Lysostaphin and N-acetylmuramidase were enhanced in comparison with those by the single enzyme.

Treatment temperature with enzymes on *Staphylococcus aureus* should preferably be adjusted to from about 4°C to about 60°C, then that on *Staphylococcus epidermidis* should be adjusted to about 25°C or more, preferably, to about 37°C or more, and that on *Enterococcus faecalis* should be adjusted to from about 25°C to less than about 60°C, preferably, from about 37°C to about 42°C. Accordingly, it is most preferable to designate the temperature range of from about 37°C to about 42°C as the optimum treatment temperature. Further, the critical temperature range to be shared with the three bacterial species is expected to the range of from about 26°C to about 59°C.

Then, treatment time with enzymes on any digested sample from *Staphylococcus aureus*, *Staphylococcus epidermidis*, *Enterococcus faecalis* is 20 minutes or more (both at zero minute and 10 minutes were not appropriate), then, since there was no leukocyte in the bacteria, the treatment time should be adjusted to at least about 15 minutes or more, preferably, to about 20 minutes or more, and be designated the time range of from about 30 minutes to about 60 minutes as the optimum treatment time. Treatment time of from about 15 minutes to about 120 minutes may also be applicable.

Then, N-acetylmuramidase is an enzyme which lower an absorbance at 600nm under the reaction at 37°C for 5 minutes of N-acetylmuramidase with heat-treated dry-powder of *Enterococcus faecalis* in 5mmol/l Tris-HCl Buffer Solution (pH

6.0) containing 2 mmol/l magnesium chloride. Then, when 1 Unit of enzyme activity is defined as an activity to lyse one ug of heat-treated dry-powder of *Streptococcus salivarius* (IFO 3350) in one minute which was determined thereon at 37°C and pH 7.0, the enzyme of 2,000 Units/mg or more is preferable.

Lysozyme is an enzyme which lower an absorbance at 600nm under the reaction of *Micrococcus luteus* with Lysozyme in PBS at 37°C for 5 minutes. Then, when 1 Unit of enzyme activity is defined as an activity to lower 0.001 of an absorbance at 540nm in one minute which was determined on *Micrococcus luteus* at 35°C and pH 6.2, an enzyme of 50,000 Units/mg or more is preferable.

Lysostaphin is an enzyme which lower an absorbance at 600nm under the reaction of *Staphylococcus epidermidis* with Lysostaphin in PBS at 37°C for 5 minutes. When 1 Unit of enzyme activity is defined as an activity to lower an absorbance at 620nm of 0.240 to 0.125 in 10 minutes which was determined on *Staphylococcus aureus* at 37°C and pH 7.5, an enzyme of 500 Units/mg or more is preferable.

Zymolase (Products Name: Zymolyase (SEIKAGAKU CORPORATION)) is an enzyme taken from the liquid culture medium of *Arthrobacter lutesul* and have strong degradation activities on cell walls of the active yeast cells. Essential enzymes contained in Zymolase and involved with cell wall degradation is  $\beta$ -1,3-glucan laminaripentaohydrolase which acts on glucose polymer with  $\beta$ -1,3-bonds and produces laminaripentaose as a main product. Zymolyase-100T is purified in ammonium sulfate fractionation, then in affinity chromatography (Kitamura, K. et al., *J. Ferment. Technol.*, **60**, 257, 1982) and have activity of 100,000 Units/g. However, it is well known that activities of the subjected enzyme are changed according to the kinds of yeast to be substrates, culture condition and growth phase thereof (Kitamura, K. et al., *J. Gen. Appl. Microbiol.*, **20**, 323, 1974; Kitamura, K. et al., *Agric. Biol. Chem.*, **45**, 1761, 1981; Kitamura, K. et al., *Agric. Biol. Chem.*, **46**, 553, 1982).

Zymolyase-100T contains about  $1.0 \times 10^7$  Units/g of  $\beta$ -1,3-glucanase, about  $1.7 \times 10^4$  Units/g of protease and about  $6.0 \times 10^4$  Units/g of mannase, but does not contain any DNase and RNase (Kitamura, K. et al. ; *J. Gen. Appl. Micro-biol.*, 18, 57, 1972). Then, the optimum pH of Zymolyase is from about 5.5 to about 8.5, preferably, from about 6.5 to about 7.5, while the optimum temperature thereof is from about 25°C to about 55°C, preferably, from about 35°C to about 45°C. Further, lytic spectrum (genus) to yeast (cells in logarithmic growth phase) may includes *Ashbya*, *Candida*, *Debaryomyces*, *Eremothecium*, *Endomyces*, *Hansenula*, *Hanseniaspora*, *Kloeckera*, *Kluyveromyces*, *Lipomyces*, *Helschkowia*, *Pichia*, *Pullularia*, *Torulopsis*, *Saccharomyces*, *Saccharomycopsis*, *Saccharomycodes*, *Schwanniomyces* or the like.

In particular, *Candida* genus may include *Candida albicans*, *Candida tropicalis*, *Candida parasilosis*, *Candida galacta*, *Candida guilliermondii*, *Candida krusei*, *Cryptococcus neoformans*. SH compounds, for example, cysteine, 2-mercapto-ethanol, dithiothreitol can be employed as an activator of these enzymes.

Bacteria belonged to these genus may also be employed in the present invention. 1 Unit of the enzyme activity is defined as an activity to lower about 30% an absorbance at 800nm of the reaction solution (prepared by adjusting with 1 ml of sterilized-purified water to be 10 ml as their final volume containing 1 ml enzyme solution of 0.05~0.1 mg/ml, 3ml of Brewer's Yeast Suspension (2 mg dry-weight/ml) as a substrate, and 5 ml of M/15 Phosphoric Acid Buffer Solution (pH 7.5)) in two hours which was determined on Brewer's Yeast Suspension as a substrate at about 25°C. Zymolyase-100T has an activity of 100,000 Units/g.

With regard to the concentration of PMSF (to be added to keep a form of the leukocytes by protecting them from protease) to be used as a reagent solvent, since 10  $\mu$ mol/l or more of PMSF

concentration was effective and a form of the leukocyte was completely kept at the PMSF concentration of 0.1 mmol/l or more, PMSF concentration should be adjusted to the range of from about 10  $\mu$ mol/l to about 10 mmol/l, preferably that of from about 0.1 mmol/l to about 1 mmol/l. Similarly, DMSO concentration should also be adjusted to less than 5%, preferably 2% or less, and more preferably about 1%. Accordingly, the reagent solvent should preferably be prepared by diluting 100 to 1,000-fold, with PBS, Dimethyl Sulfoxide (DMSO) containing 0.1 mol/l Phenylmethyl Sulfonylfluoride (PMSF).

Acetylation of cell membrane proteins may further be performed after the step to bare DNA in the causative microorganisms. In particular, slides are immersed in the acetylation reagent prepared by adding Acetic Anhydride to the acetylation raw reagent (adjusted with quantum sufficient of sterilized-purified water to be 50ml as their total volume containing 7.46g of Triethanolamine and quantum sufficient of Hydrochloric Acid), diluting from about two-fold to about 50-fold, preferably about 10-fold with sterilized-purified water, and adjusting final concentration of Acetic Anhydride to from 0.1% to 3.0%, preferably 0.8%, and are applied to a centrifuge for from 5 to 30 minutes. After then, they are completely air-dried by immersing them successively into 75%, 85% and 98% Ethanol for from two to five minutes respectively.

Alkalization of DNA in the causative microorganisms may further be performed to reshape it into the single strand DNA after the step to acetylate the cell membrane proteins. In particular, slides are immersed for from about two minutes to about five minutes in PBS containing from about 10 mmol/l to about 300 mmol/l Sodium Hydroxide, preferably, about 70 mmol/l Sodium Hydroxide (prepared by adding Sodium Hydroxide to PBS Raw Solution, diluting 20-fold with sterilized-purified water, and adjusting final concentration of Sodium Hydroxide to 70 mmol/l). After then, they were completely air-dried by immersing them successively into 75%, 85% and 98% Ethanol for

from two to five minutes respectively.

In order to perform *in situ* hybridization with detective DNA probe which can hybridize under the stringent condition to the naked DNA from the causative microorganisms, probe solution (solution containing the detective DNA probe prepared with probe dilution) is applied to the area to be smeared, and is left for from about one hour to about three hours, preferably, for about two hours in the wet chamber of from about 25°C to about 50°C, preferably, from about 37°C to about 42°C.

Then, three stained bottles containing the hybridization detergent (prepared by mixing Hybridization Raw Solution (prepared with sterilized-purified water to be 75ml as their total volume containing 13.15g of Sodium Chloride, 6.615g of Trisodium Citrate Dihydrate) in the ratio of Hybridization Raw Solution : sterilized-purified water : Formamide = 5:45:50) are provided and they are successively immersed therein at from about 35 to about 45°C, preferably, at about 42°C for 10 minutes respectively. They are then immersed in PBS and are applied successively to centrifugation on a centrifuge for from about five to about 30 minutes. In particular, the diluted probe solution contains 600  $\mu$ l of Salmon Sperm DNA, 50  $\mu$ l of 100 $\times$  Denhert's Solution, 500  $\mu$ l of Hybridization Raw Solution, 2250  $\mu$ l of Formamide and 1000  $\mu$ l of 50% Dextran Sulfate. Probe solution contains preferably 15ng of each detective DNA probes and may be adjusted with the probe dilution to be 50  $\mu$ l as their total volume.

Probe concentration of SA, SE, PA, EF, EK is adjusted to from about 0.6 ng/ $\mu$ l to about 1.8 ng/ $\mu$ l, preferably to from about 0.6 ng/ $\mu$ l to about 1.2 ng/ $\mu$ l. Since result at 0.06 ng/ $\mu$ l was not acceptable while result at 0.6 ng/ $\mu$ l was acceptable, it is preferably to adjust the concentration to at least 0.1 ng/ $\mu$ l or more. Then, since result at 2.4 ng/ $\mu$ l was not acceptable while result at 1.8 ng/ $\mu$ l was acceptable, it is preferably to adjust the concentration to 2.2 ng/ $\mu$ l or less. Further, the optimum concentration of the positive control and



the negative control are adjusted to concentration of from 0.4 ng/ $\mu$ l to 2.0 ng/ $\mu$ l and from 0.6 ng/ $\mu$ l to 2.0 ng/ $\mu$ l respectively, preferably, that of from 0.6 ng/ $\mu$ l to 1.0 ng/ $\mu$ l for the both control.

Time to perform hybridization is at least 30 minutes or more, preferably 60 minutes or more, and more preferably 90 minutes or more. Time of from about 120 minutes to about 900 minutes may be designated as optimum hybridization time.

It is preferably to use surfactants like sodium dodecyl sulfate (SDS) in the step of *in situ* hybridization because it may allow enhancement of the detection sensitivity. SDS concentration is preferably 1% or less, more preferably from about 0.1% to about 0.5%, more further preferably about 0.25%. SDS is added to the solution to be employed at the hybridization, otherwise, it may be added in advance to the probe diluent or the probe solution.

It is recommended to employ, as the detective DNA probes, one or more kind(s) of DNA probe(s) having from about 350 to about 600 base length, preferably from about 350 to about 550 base length, because the probes are thereby smoothly incorporated into the phagocytes and they may easily and exactly be contacted with genes of the exogenous microorganisms wherein such probes are incorporated. It is not necessary to fall base length (number of bases) of the subjected within the base length range aforementioned, but is simply recommended to employ probes having base length distribution to be at least partially overlapped with the base length range aforementioned. These probes are made of single or several (one or more) kind(s) of probes. One or more probe(s) may be plural kinds of probes to be hybridized to the single bacterial species or may be plural kinds of probes to be hybridized to the plural bacterial species one to one, but any restriction would not be imposed as far as one or more kind(s) of probes is/are used.

These probes have preferably DNA fragments containing a sequence not to be hybridized anyway to phagocytes themselves and would not cross-hybridize at all to any gene from the irrelevant bacteria species. Specific probes would be prepared in a short time, for example, with a subtraction method. These probes may be prepared and labelled through the conventional nick translation methodology using non-radioisotopic labelling substances like fluorescein isothiocyanate (FITC), biotin, digoxigenin (digoxigenin(DIG)-11-dUTP) or the like. Strand length of the probes can be adjusted most effectively by changing the amount ratio of DNaseI and DNA polymerase I respectively to be added at the nick translation reaction. For example, in order to effectively label 2  $\mu$ g of DNA probe (SA-24) and adjust strand length thereof (into base length of from about 350 to about 600) which allow the probes to be effectively *in situ* hybridized to the exogenous microorganisms DNA, in the reactionary solution of 100  $\mu$ l total volume, with regard to 2  $\mu$ l of 10U/ $\mu$ l DNA polymerase I, 6  $\mu$ l of DNaseI is presented in the 100  $\mu$ l total volume as an activity of from about 10 mU to about 350 mU, preferably from about 25 mU to about 200 mU, and more preferably from about 50 mU to about 150 mU. As far as volume ratio in the essential optimum reaction conditions aforementioned is kept at the certain level, volume of each enzyme and total volume of the reactionary solution can be changed optionally. In the other words, volume of DNaseI may be adjusted, with regard to 20 U of DNA polymerase I in the total volume of 100  $\mu$ l, to from about 10 mU to about 350 mU, preferably from about 25 mU to about 200mU, and more preferably from about 50 mU to about 150 mU. Turning to the other aspect, it is recommended to perform nick translation reaction by adjusting volume of DNaseI, with regard to 1 Unit of DNA polymerase I, to from about 0.5/1,000 U to about 17.5/1,000 U, preferably from about 1.25/1,000 U to about 10/1,000 U, and more preferably from about 2.5/1,000 U to about 7.5/1,000 U. Then, with regard to 1  $\mu$ g of DNA, volume of DNA polymerase I is adjusted to about 10 U, while that of DNaseI is adjusted to from about 5 mU to about 175mU, preferably from about 12.5 mU to about 100 mU, and more preferably from

about 25 mU to about 75mU. With regard to the other probes, an amount of DNA as well as optimum reaction conditions on DNA polymerase and DNase I are determined by referring to the optimum reaction conditions aforementioned, then the probe length (the length of from about 350 bases to about 600 bases) is also determined in which the length may allow the probes to be effectively labelled and may allow the probes to be effectively *in situ* hybridized to the exogenous microorganisms DNA.

Stringent condition to be employed at the *in situ* hybridization is a condition, for example, to incubate under the presence of formamide of from about 30% to about 60%, preferably at about 50%, at from about 30 °C to about 50 °C, preferably from about 38°C to about 42°C, and rinse successively.

After *in situ* hybridization, blocking operation may also be performed. In particular, 1 ml of Blocking reagent (prepared with sterilized-purified water to be 10ml as their total volume containing 2 ml of Rabbit Normal Serum and 0.5ml of PBS Raw Solution) per single slide is dropped onto the smeared area thereof and the slides were left for from 15 to 60 minutes. Then, Blocking reagent is removed.

In order to detect the signals resulted from hybridization with the bacterial gene (genome DNA or RNA), any coloration reaction utilizing the conventional antigen-antibody reaction or the like may be employed. Namely, the samples so hybridized are fully washed, then are subjected to Blocking, and are treated with complexes like anti-FITC antibody, anti-digoxigenin antibody, for example, alkaline phosphatase complexes, followed by evaluation on the hybridization results through signals to be expressed by the chromogenic substrates in the complexes. For example, when a probe is labelled with digoxigenin 11-dUTP as noted above, anti-digoxigenin alkaline phosphatase complexes will be employed, and the probe may be detected with a substrate (Nitro Blue Tetrazolium, 5-Bromo-4-Chloro-3-Indolyl Phosphate or the like) to be usually employed on the alkaline phosphatase.

Then, the smear samples, which are washed after the coloration reaction, are subjected to counterstain with Naphtol Black, Fast Green (20mg/50ml, Wako Chemicals) or the like, and in-cell signals are detected by an optical microscopy.

Particularly, in order to take signals in hybridization by employing, for example, DNA probes labelled with digoxigenin as detective DNA probes, labelled antibody solution is prepared by diluting from 10 to 200-fold, preferably 50-fold the labelled antibody (prepared with 12.6  $\mu$ l of Buffer A (prepared with quantum sufficient of sterilized-purified water to be 100 ml as their total volume containing 746 mg of Triethanolamine, 17.5 mg of Sodium Chloride, 20.3 mg of Magnesium Chloride Hexahydrate, 1.36 mg of Zinc Chloride, 1000 mg of Bovine Serum Albumin and quantum sufficient of Hydrochloric Acid) to be 14  $\mu$ l as their total volume containing 1.05 Unit of alkaline-phosphatase-labelled anti-digoxigenin antibody solution) with the antibody dilution (prepared with quantum sufficient of sterilized-purified water to be 0.7ml as their total volume containing 8.48 mg of Tris-(Hydroxymethyl)-Aminomethane, 6.14 mg of Sodium Chloride, and quantum sufficient of Hydrochloric Acid), then 10  $\mu$ l of the labelled antibody solution is dropped onto each of the smeared area, and they may be left for from 15 to 60 minutes. After then, they are immersed in from two-fold to fifty-fold diluted solution, preferably ten-fold diluted solution of the labelled antibody detergent solution (prepared with sterilized-purified water to be 100ml as their total volume containing 1ml of Polysolvate 20 and 50ml of PBS Raw Solution) and are centrifuged for from five to 30 minutes during which they are being immersed in the detergent solution. This operation is repeated twice, then the samples are immersed in the treatment solution prepared by mixing Preliminary Treatment Solution 1 (prepared with quantum sufficient of sterilized-purified water to be 50ml as their total volume containing 6.06 g of Tris-(Hydroxymethyl)-Aminomethane, 2.92 g of Sodium Chloride, and quantum sufficient of Hydrochloric Acid) with equal amount of Preliminary Treatment Solution 2 (prepared with quantum

sufficient of sterilized-purified water to be 50ml as their total volume containing 5.08 g of Magnesium Chloride Hexahydrate), then diluting approximately 5-fold with sterilized-purified water, and are centrifuged for from five to 30 minutes during which they are being immersed in the detergent solution. 1 ml of chromogenic agent (Nitro Blue Tetrazolium (NBT)/ 5-Bromo-4-Chloro-3-Indolyl Phosphate (BCIP) per single slide is dropped onto the smeared area of the slides by filtrating with a disposable syringe equipped with 0.2  $\mu$ m syringe top filter, and is kept in the dark and is left in a wet chamber for from about 15 minutes to about 60 minutes at from about 10°C to about 45°C, preferably at 37°C. Then, they are immersed for from about two to about 10 minutes in the solution prepared by diluting from two-fold to about 50-fold, preferably about ten-fold the chromogenic agent cleaner (prepared with quantum sufficient of sterilized-purified water to be 50ml as their total volume containing 606mg of Tris-(Hydroxymethyl)-Aminomethane, quantum sufficient of Hydrochloric Acid and 186mg of Disodium Ethylenediaminetetraacetic acid Dihydrate) and are air-dried. Further, they are immersed in the solution prepared by diluting from two-fold to 50-fold, preferably 10-fold the counterstain solution (prepared with quantum sufficient of sterilized-purified water to be 50ml as their total volume containing 50mg of Fast Green FCF (Edible Dye Green No.3)) and in an acetic acid solution of from about 0.1% to about 5%, preferably about 1%. Thereafter, they may be immersed again in the solution prepared by diluting about two-fold to about 50-fold, preferably 10-fold the cleaner aforementioned to remove excessive counterstain solution and are completely air-dried. Each of such chromogenic agents may be prepared individually.

Preferable solution of the anti-digoxigenin antibodies labelled with alkaline phosphatase may include a solution which will offer color in the blotted area made by blotting one ng of DNA labelled with digoxigenin onto a blotting membrane, blocking the same, then treating those with the 10,000-fold diluted solution of the anti-digoxigenin antibody labelled with

alkaline phosphatase, and reacting those with chromogenic substrates (NBT/BCIP), while it will not offer any color according to the same procedure employing DNA without digoxigenin label. Anti-digoxigenin antibodies taken from sheep are preferable. In particular, it is recommended to take such antibodies by purifying the immunized sheep serum with an ion-exchange chromatography and an antibody column chromatography.

Chromogenic agent (NBT/BCIP solution, pH 9.0-10.0) is an agent prepared preferably with quantum sufficient of sterilized-purified water to be 10ml as their total volume containing 3.3mg of Nitro Blue Tetrazolium (NBT), 1.65mg of 5-Bromo-4-Chloro-3-Indolyl Phosphate (BCIP), 99  $\mu$ g of N,N-dimethyl-formamide, 121mg of Tris-(Hydroxymethyl)-Aminomethane, quantum sufficient of Hydrochloric Acid, 58.4mg of Sodium Chloride, 101.6mg of Magnesium Chloride Hexahydrate.

Preferable chromogenic agents may include an agent which will offer violet signals in the blotted area made by blotting proteins labelled with alkaline phosphatase onto a blotting membrane and treating the same with such chromogenic agents in the dark at room temperature.

In the counterstain above, edible dye, for example, Yellow No.4 (Tartrazine) can be used to present more clearly contrast between signals and cells. Factors of such poor countercontrast may include similarity of the colors to be expressed, namely, between violet color offered by the substrate and blue color offered by Naphtol Black. When this methodology is applied to the present invention, it came to know that such methodology is useful at the counterstain. Any conventional methodology has never employed a food dye.

Nick translation methodology is applicable as a method to label the digoxigenin. The other methodology may include, for example, PCR Method, Random Primer Labelling Method, *in vitro*

Transcription Labelling Method, Terminal Transferase Labelling Method or the like.

When at least one expressed violet signal(s) is/are confirmed by the optical microscopy ( $\times 1,000$ ) on the subjected cells in the single well which is stained with the counterstain solution aforementioned, the sample may be designated as positive.

Then, Japanese Patent Nos. 2558420, 2798499, 2965543, 2965544 and 3026789 can be referred at the producing of the detective probes.

For example, in order to culture bacteria taken from working-cell-banks, the working-cell-banks (SA-24) are smeared streakily with a platinum loop, a disposable plastic loop or the like onto L-broth solid medium containing  $50 \mu\text{g/ml}$  ampicillin mounted on the sterilized laboratory dishes (Screening).

After an overnight cultivation thereof, single colony so cultured is inoculated into 5 ml of L-broth medium containing  $50 \mu\text{g/ml}$  ampicillin and is shaking cultured overnight at  $37^\circ\text{C}$  (Precultivation).

2.5 ml of the culture solution is inoculated individually into 400ml of the medium in a flask and is shaking cultured overnight at about  $37^\circ\text{C}$  (Main Cultivation).

Then, in order to extract SA-24 plasmid DNA, the cultured solution prepared through the main cultivation are centrifugated at  $4^\circ\text{C}$ , in  $4,000 \times g$ , for 10 minutes to collect the bacteria. Supernatant are removed, 20ml of STE (10 mmol/l Tris-HCl (pH 8.0), 1 mmol/l Disodium Ethylenediaminetetraacetic acid (EDTA), 0.1 mmol/l Sodium Chloride) are added to the remained bacteria to resuspend the same, and they are centrifugated at  $4^\circ\text{C}$ , in  $4,000 \times g$ , for 10 minutes to collect the bacteria. The bacteria is suspended with 5 ml of Solution-1 (50 mmol/l glucose, 25 mmol/l

Tris-HCl (pH 8.0), 10 mmol/l EDTA) containing 10mg/ml Lysozyme and is left for five minutes under the room temperature. Thereafter, 10 ml of Solution-2 (0.2 mmol/l Sodium Hydroxide, 1% Sodium Dodecyl Sulfate (SDS)) is added, then they are upset to mix, and are left for ten minutes on ice. 7.5 ml of Iced Solution-3 (3 mol/l potassium acetate (pH 4.8)) is added, then they are upset to mix, and are left for ten minutes on ice.

After centrifugation by a high speed refrigerated centrifuge at 4°C for 30 minutes at 45,000×g, the supernatant is recovered, and left to stand to cool to room temperature. After leaving to stand, 0.6 volume of isopropanol (about 24 ml) is added thereto, mixed by inversion and left to stand at room temperature for 5 minutes or longer. After centrifugation by a high speed refrigerated centrifuge at 25°C, for 30 minutes at 28,000×g, the supernatant is discarded, and thus resulting pellet is washed with 70% ethanol and air dried. After air drying, 8 ml of TE (10 mmol/l Tris-hydrochloric acid (pH 8.0), 1 mmol/l EDTA) is added thereto to dissolve the pellet (extraction of plasmid DNA).

Next, for the purification of the plasmid DNA containing SA-24, 800  $\mu$ l of 10 mg/ml ethidium bromide and 8.6 g of cesium chloride are added to the resulting plasmid DNA followed by mixing by inversion to dissolve the plasmid. The solution is placed in a centrifuge tube, which is then capped or sealed. After centrifugation at 20°C for 5 hours at 500,000×g with a vertical rotor, a band of the plasmid DNA is fractionated using a glass syringe or an injection needle under the irradiation of an ultraviolet ray light. To the fractionated plasmid DNA solution is added an equivalent amount of TE-saturated 1-butanol followed by mixing by inversion and centrifugation at 15,000 ×g for 5 minutes by a high speed microcentrifuge to remove the supernatant. This operation is repeated to eliminate ethidium bromide in the plasmid DNA solution. Next, thereto is added TE to give the volume of 1.5 ml followed by desalting on a demineralization column (NAP-10). To the desalted plasmid DNA



solution is added 30  $\mu$ l of a 3 mol/l sodium acetate solution followed by mixing, and 3 fold amount of 99.5% ethanol is added thereto followed by mixing by inversion and leaving to stand at -20°C for 30 minutes or longer. After leaving to stand, centrifugation is conducted with a high speed refrigerated micro centrifuge at 4°C for 20 minutes at 15,000 $\times$ g to remove the supernatant. Thereafter, cold 70% ethanol is added thereto to suspend therein, and once again, centrifugation is conducted with a high speed refrigerated micro centrifuge at 4°C for 20 minutes at 15,000 $\times$ g to remove the supernatant. Thus resulting precipitate of the plasmid DNA is evaporated to dryness under a reduced pressure. TE in an amount of 100  $\mu$ l is added to the plasmid DNA to dissolve completely, and the concentration is measured on the basis of the absorbance at 260 nm (Purification of plasmid DNA containing SA-24). Then, size check of the plasmid DNA containing SA-24 is carried out by a treatment with arestriction enzyme and agarose electrophoresis.

In order to purify SA-24 by digesting the plasmid DNA containing SA-24 with restriction enzyme(s) and successively applying them to an agarose electrophoresis, 1 mg of the plasmid DNA with SA-24, molecular weight of which had been determined, are digested through a reaction to be performed at 37°C for one and half hours or more under the presence of the restriction enzyme HindIII alone or the combination of HindIII and the other restriction enzyme(s). After digesting the plasmid DNA, completion of such digestion is confirmed by applying a part of the reactionary solution to 0.8% agarose electrophoresis. Then, after confirming such digestion, SA-24 band is taken by applying the solution to an electrophoresis with 0.8% agarose as a fractionator. SA-24 so taken are extracted from the agarose gel and are purified, then, concentration thereof is determined with an absorptiometer. A part of the purified SA-24 is applied to an electrophoresis with 0.8% agarose gel and is appeared as a single band.

SA-24 may preferably be labelled with digoxigenin with the

reactionary solution having the composition listed in the following Table 1 including  $2\mu\text{g}$  of the purified SA-24.

**TABLE 1**

Composition in Reactionary Solution to Label

	Amount ( $\mu\text{L}$ )
DNA probe	X
$10\times\text{L.B.}^{(a)}$	10
100mmol/L dithiothreitol	10
dNTPs <sup>(b)</sup> (A, G, C 0.5mmol/L)	4
digoxigenin-dUTP <sup>(c)</sup> (0.4mmol/L)	5
DNaseI <sup>(d)</sup>	6
10U/ $\mu\text{L}$ DNA polymerase I	2
Sterilized-Purified Water	Y
Total	100

[REMARKS]

- (a)  $10\times\text{L.B.}$  : 0.5 mol/L Tris-HCl (pH 7.5),  
50 mmol/L magnesium chloride,  
0.5 mg/mL bovine serum albumin
- (b) dNTPs : 0.5 mmol/L 2'-dioxadenosine-5'-triphosphate,  
0.5 mmol/L 2'-dioxycytidine-5'-triphosphate,  
0.5 mmol/L 2'-dioxycytidine-5'-triphosphate
- (c) digoxigenin-dUTP : 0.4mmol/L digoxigenin-11-2'-  
dioxycytidine-5'-triphosphate
- (d) DNase I : Deoxyribonuclease I is diluted with 25 mmol/L  
Tris-HCl (pH 7.5), 50% glycerin solution to be  
used at 50-150 mU per total amount of  $100\mu\text{L}$  and  
is adjusted to the amount aforementioned.

In Table 1, the volume X is adjustable to realize preferable probe concentration aforementioned according to the concentration of raw probe solution, then, the final volume is adjusted by determining, based on the volume X, the volume Y of Sterilized- Purified Water. After putting the labels, the

reaction is terminated by adding 100  $\mu$ l of TE to the reaction solution.

Free nucleotides are then removed by adding solution to terminate the reaction into the spin column and centrifuging the same at 4  $^{\circ}$ C, in 380 $\times$ g, for 10 minutes. Thereafter, concentration of the eluted solution is determined with an absorptiometer and is adjusted with TE to the unit level of ng/ $\mu$ l.

To confirm the labelled subjects, 0.5  $\mu$ l of the labelled SA-24 is dropped onto a membrane and is air-dried. The membrane is immersed into the blocking reagent and is kept for 30 minutes under the room temperature. The membrane is then immersed for 30 minutes under the room temperature into the solution of anti-digoxigenin antibody labelled with alkaline phosphatase prepared by diluting 5,000-fold with a solution containing 0.1mol/l Tris-HCl (pH 7.5) and 0.15 mol/l sodium chloride. The membrane is immersed into a solution containing 0.1 mol/l Tris-HCl (pH 7.5) and 0.15 mol/l sodium chloride, then is shaken for 10 minutes under the room temperature, and is rinsed twice. Further, the membrane is immersed for 10 minutes under the room temperature into a solution containing 0.5 mol/l Tris-HCl (pH 9.5), 0.15 mol/l sodium chloride and 50 mmol/l magnesium chloride. The membrane is then immersed into chromogenic agent for 10 minutes, in the dark, under the room temperature. Labelled subjects are confirmed with violet color appeared under the spotted position.

Spin column is produced by filling a few amount of sterilized grass wool into 1 ml volume of disposable syringe. Sephadex G-50 swelled with 1 mmol/l Tris-HCl (pH 7.5), 1 mmol/l EDTA and 0.1% SDS is then filled into the syringe. The syringe is inserted into 15ml volume of the disposable conical tube and is centrifuged at 4 $^{\circ}$ C, in 320 $\times$ g, for 10 minutes to remove the excess buffer solution. The syringe is then detached from the disposable conical tube and the excess buffer solution is discarded. Thereafter, the spin column is assembled by engaging 1.5ml volume of Eppendorf Tube with the bottom of the disposable

conical tube and entering the syringe into the conical tube.

Dot Blot Hybridization according to the following procedure is recommended to confirm specificity of the probes.

First of all, in order to denature each of the spotted genomic DNA, according to the standard procedure, 100ng of the various bacterial genome so prepared are spotted onto nylon membranes (Pall BioDine Type B; Nihon Pall Ltd.), and air-dried membranes are kept for 10 minutes on the filter papers (3mm: Wattman) saturated with solution containing 0.5mol/l sodium hydroxide and 1.5mol/l sodium chloride. The denatured DNA are then neutralized by leaving them for 10 minutes on the previously noted filter papers saturated with solution containing 0.5mol/l Tris-HCl (pH 7.5) and 1.5 mol/l sodium chloride. They are further left for five minutes on the previously noted filter papers saturated with  $2 \times$  SSC (Standard Saline Citrate) Solution and are rinsed. After then, such membranes are air-dried and are immersed into  $2 \times$ SSC Solution for five minutes. In accordance with the standard procedure, the membranes are immersed into Prehybridization Solution in plastic bags and are kept at 42°C for 60 minutes. In the plastic bag, the membranes are immersed in 15 ml of Hybridization Solution containing 400ng of Probes and are reacted overnight at 42°C. Next, the membranes are immersed into the solution containing  $2 \times$ SSC and 0.1% SDS (sodium dodecyl sulfate) and are rinsed for five minutes (this is repeated twice). The membranes are then immersed into the solution containing 0.1 $\times$ SSC, 0.1% SDS and are rinsed for ten minutes (this is repeated three times). The membranes are immersed into  $2 \times$ SSC solution and are rinsed for five minutes. The membranes are immersed into the solution containing 3% bovine serum albumin, 1% Blocking Buffer (Boehringer), 0.1 mol/l Tris-HCl (pH 7.5) and 0.15mol/l Sodium Chloride, and are gently shaken for 30 minutes. The membranes are immersed into the solution prepared by diluting 5,000-fold the alkaline-phosphatase labelled anti-digoxigenin antibody (Boehringer) with the solution containing 0.1 mol/l Tris-HCl (pH 7.5) and 0.15 mol/l Sodium Chloride, and are gently shaken for 30 minutes.

Next, the membranes are immersed into the solution containing 0.1 mol/l Tris-HCl (pH 7.5) and 0.15 mol/l sodium chloride, and are shaken for 15 minutes (twice). The membranes are immersed into the solution containing 0.1 mol/l Tris-HCl (pH 9.5), 0.1 mol/l Sodium Chloride and 5 mmol/l Magnesium Chloride, and are shaken for 5 minutes. The membranes are immersed into NBT-BCIP Solution (GIBCO BRL) and are subjected to chromogenic reaction in the dark. The membranes are then immersed into TE (10 mmol/l Tris-HCl (pH 8.0), 1 mmol/l EDTA) to terminate the chromogenic reaction and are air-dried. Particulars of Prehybridization Solution and Hybridization Solution are illustrated in the following Table 2.

TABLE 2

[Unit : ml]

	Prehybridization Solution	Hybridization Solution
Formamide	7. 5	6. 7 5
20×SSC Solution	3. 7 5	3. 7 5
100×Denhart Solution	0. 7 5	0. 1 5
0.5mol/L Phosphatized Buffer Solution	0. 7 5	0. 6
Sterized-Distillated Water	1. 5	1. 9 5
10mg/mL Salmon Sperm DNA	0. 7 5	0. 3
50% Dextran Sulfate	— — — —	1. 5
Total Liquid Amount	1 5. 0	1 5. 0

Conventional surfactants can be employed as those to be used in the step of *in situ* hybridization. Typically, the surfactants are roughly divided into anionic surfactant, non-ionic surfactant, cationic surfactant and amphoteric surfactant.

Anion surfactants are also referred to as anionic surfactants, which yield an organic anion upon ionization in water. When a lipophilic group in the molecule of the surfactant is represented by R, examples of the anion surfactant include  $\text{RCOONa}$ ,  $\text{RSO}_3\text{Na}$ ,  $\text{RSO}_4\text{Na}$  and the like. An aqueous solution of the surfactant containing a weakly acidic group such as  $\text{RCOONa}$  is liable to be hydrolyzed and is weak alkaline. However, an aqueous solution of a surfactant having a strongly acidic group such as  $\text{RSO}_3\text{Na}$ ,  $\text{RSO}_4\text{Na}$  or the like is resistant to hydrolysis, which shall be neutral. Because it is anionic, it may lose surface activity in the presence of a large quantity of cationic substance, and may be inactivated in a strongly acidic circumstance.

Nonionic surfactants refer to those having a hydrophilic group which is nonionic. An ethylene oxide group ( $-\text{CH}_2\text{CH}_2\text{O}-$ ) is often used as the hydrophilic group. As number of this group increases, hydrophilicity is increased. To the contrary, as number of the lipophilic group increases, lipophilicity is increased. Therefore, it is characterized in that surfactants with variously altered hydrophilicity and lipophilicity can be obtained. Because a nonionic surfactant does not ionize in water and is hardly affected by inorganic salts, less action is exerted also on a living body. In addition, the detergent action thereof is potent with comparatively less foaming, therefore, it is widely used not alone as a detergent, but in pharmaceuticals, cosmetics, foods and the like. Water soluble nonionic surfactant becomes insoluble in water at a certain temperature as the temperature rises, and then the aqueous solution starts to be turbid. Such turbidity results from the cleavage of hydrogen bonds between the hydrophilic groups and water.

Cation surfactants are also referred to as cationic surfactants, which yield an organic cation upon ionization in water. Although cation surfactants do not have potent detergent action in general, they strongly bind to anionic substances such

as bacteria, leading to a great bactericidal action. Moreover, they also have an anti-static ability for fibers and plastics. Although dodecyltrimethyl chloride  $[C_{12}H_{25}(CH_3)_3N]Cl$  as a typical exemplary cation surfactant is water soluble, didodecyldimethylammonium chloride  $[(C_{12}H_{25})_2(CH_3)_2N]Cl$  is insoluble in water, which forms a vesicle in the form of a bimolecular film in water, and is soluble in benzene.

Ampholytic surfactants are surfactants having both an anionic group and a cationic group in the molecule. Ionization state thereof in water is similar to those of amino acids, and thus many of ampholytic surfactants are amino acid derivatives. Therefore, they have an isoelectric point similarly to amino acids, which act as an anion surfactant in an alkaline region from the isoelectric point, whilst as a cation surfactant in an acidic region. Water solubility becomes the lowest at the isoelectric point, and the surface tension is also reduced. Ampholytic surfactants are used for a bactericide, an antistatic agent or the like.

Furthermore, anion surfactants are classified into the carboxylic acid type, sulfonic acid type, sulfate ester type and phosphate ester type, whilst nonionic surfactants are classified into the ester type, ether type, ester ether type and alkanolamide type. Cation surfactants are classified into alkylamine salt type and quaternary ammonium salt type, whilst ampholytic surfactants are classified into carboxy betaine type, 2-alkylimidazoline derivative type and glycine type.

Moreover, the anion surfactants of carboxylic acid type are further classified into fatty acid monocarboxylate salts, N-acylsarcosine salts and N-acylglutamate salts. Representative examples thereof respectively include: sodium laurate and medicated soap as the fatty acid monocarboxylate salts; sodium N-lauroylsarcosine as the N-acylsarcosine salt; and disodium N-lauroylglutamate as the N-acylglutamate. Still more, the sulfonic acid type is further classified into dialkyl



sulfosuccinate salts, alkane sulfonate salts, alpha-olefin sulfonate salts, straight chain alkyl benzenesulfonate salts, alkyl (branched chain) benzenesulfonate salts, alkyl naphthalenesulfonate salts, naphthalenesulfonate salts-formaldehyde condensates and N-methyl-N-acyltaurine salts. Representative examples include: sodium dioctyl sulfosuccinate as the dialkyl sulfosuccinate salt; sodium dodecane sulfonate as the alkane sulfonate; sodium straight chain dodecyl benzenesulfonate as the straight chain alkyl benzenesulfonate salt; sodium dodecyl benzenesulfonate as the alkyl (branched chain) benzenesulfonate salt; sodium butyl naphthalenesulfonate as the alkyl naphthalenesulfonate salt; and sodium N-methyl-N-stearoyltaurine as the N-methyl N-acyltaurine salt. In addition, the sulfate ester type is further classified into alkyl sulfate salts, polyoxyethylene alkyl ether sulfate salts and oil-and-fat sulfate ester salts. Representative examples include sodium dodecyl sulfate, sodium lauryl sulfate and sodium cetyl sulfate as the alkyl sulfate salt; and polyoxyethylene lauryl ether sulfate triethanolamine as the polyoxyethylene alkyl ether sulfate salt. Moreover, the phosphate ester type is further classified into alkyl phosphate salts, polyoxyethylene alkyl ether phosphate salts and polyoxyethylene alkylphenyl ether phosphate salts. Representative examples include disodium monolauryl phosphate as the alkyl phosphate salt; and sodium polyoxyethylene lauryl ether phosphate and polyoxyethylene oleyl ether phosphate (8MOL) as the polyoxyethylene alkyl ether phosphate salt.

Ester type of the nonionic surfactants is further classified into fatty acid glycerin, fatty acid sorbitan and fatty acid sucrose ester. Representative examples respectively include: glycerin monostearate as the fatty acid glycerin; sorbitan monostearate, sorbitan trioleate, sorbitan sesquioleate, sorbitan monolaurate, polysorbate 20 (polyoxyethylene sorbitan fatty acid ester), polysorbate 60 and polysorbate 80 as the fatty acid sorbitan; and stearic acid sucrose ester as the fatty acid sucrose ester. Additionally,

the ether type is further classified into polyoxyethylene alkyl ether, polyoxyethylene alkyl phenyl ether and polyoxyethylene polyoxypropylene glycol. Representative examples include: polyoxyethylene lauryl ether, polyoxyethylene stearyl ether and polyoxyethylene cetyl ether as the polyoxyethylene alkyl ether; and polyoxyethylene nonyl phenyl ether and polyoxyethylene octyl phenyl ether as the polyoxyethylene alkyl phenyl ether. In addition, the ester ether type is further classified into fatty acid polyethylene glycol and fatty acid polyoxyethylene sorbitan. Representative examples thereof respectively include oleic acid polyethylene glycol as the fatty acid polyethylene glycol; and polyoxyethylene sorbitan palmitate and polyoxyethylene sorbitan monolaurate as the fatty acid polyoxyethylene sorbitan. In addition, the alkanolamide type involves only fatty acid alkanolamide alone. Representative example is lauric diethanolamide.

The alkyl amine salt type of the cation surfactant includes monoalkyl amine salts, dialkyl amine salt and trialkyl amine salts. Representative examples thereof include monostearyl amine hydrochloride. Moreover, the quaternary ammonium salt type is further classified into alkyltrimethyl ammonium chloride (or bromide or iodide), dialkyldimethyl ammonium chloride (or bromide or iodide), and alkyl benzalkonium chloride. Representative examples respectively include: stearyltrimethyl ammonium chloride as the alkyltrimethyl ammonium chloride (or bromide or iodide); distearyldimethyl ammonium chloride as the dialkyldimethyl ammonium chloride (or bromide or iodide); and lauryl benzalkonium chloride as the alkyl benzalkonium chloride.

The carboxy betaine type of the ampholytic surfactant is only alkyl betaine alone. Representative example is lauryl betaine. Additionally, the 2-alkyl imidazoline derivative type is only 2-alkyl-N-carboxymethyl-N-hydroxyethyl imidazolinium betaine alone. Representative example includes 2-undecyl-N-carboxymethyl-N-hydroxyethyl imidazolinium betaine. In addition, the glycine type may be alkyl (or dialkyl) diethylene

triaminoacetic acid, and the representative example includes dioctyl diethylene triaminoacetic acid.

Moreover, in addition to the representative examples as described above, Triton X-100, lauryl sarcosine, saponin, BRIJ35, alkyl allyl polyether alcohol, higher alcohol sulfate, N-cocoyl-L-arginine ethyl ester DL-pyrrolidone carboxylate salt, sodium N-cocoyl-N-methyl aminoethyl sulfonate, cholesterol, self emulsifying type monostearate glycerin, squalane, stearyl alcohol, stearic acid polyoxyl 40, cetyl alcohol, cetomacrogol 1000, sebacate diethyl, nonylphenoxy polyoxyethylene ethane sulfate ester ammonium, polyoxyethylene oleylamine, polyoxyethylene sorbit yellow bees wax, polyoxyl 35 castor oil, macrogol 400, N-coconut oil fatty acid acyl L-arginine ethyl DL-pyrrolidone carboxylate salt, lauryldimethylamine oxide solution, lauromacrogol, methylcellulose, CMC (carboxymethylcellulose), polyoxyethylene hardened castor oil 20 and polyoxyethylene hardened castor oil 60, CHAPS, deoxycholic acid, digitonin, n-dodecyl maltoside, Nonidet P40, n-octyl glucoside, octyl thioglucoside, laurate sucrose, dodecyl poly(ethylene glycol ether)n, n-dodecyl-N,N-dimethyl-3-ammonio-1-propanesulfonate and the like are also included.

It is important to apply the various surfactants listed above at the *in situ* hybridization step, while application embodiment thereof is not limited anyway. For example, one skilled in the art would mix the surfactants previously into probe solution or probe dilution, otherwise, the skilled artisan may prepare the separate solution containing surfactants and add the same before, simultaneously or after application of the probe solution onto the area to be smeared.

If any positive control probe is necessary in the present invention, such probe may be prepared as follows. For example, in order to conduct the extraction and purification of the genomic DNA of U937 cell (ATCC CRL-1593.2), U937 cells are first

cultured in a 5% carbon dioxide gas incubator at 37°C using an RPMI1640 medium (25 ml) in a cell culture flask (175 cm<sup>2</sup>). The U937 culture solution is placed in a 50 ml centrifuge tube, and centrifuged at 4°C for 10 minutes at 220×g to recover the U937 cells. The cells are suspended and washed in 10 ml of PBS, and again centrifuged at 4°C for 10 minutes at 180×g to recover the cells. Thereafter, the supernatant is discarded, and the cells are suspended in 1 ml of a TE solution containing 200 µg/ml proteinase K and containing 1% SDS, followed by leaving to stand at 37°C for 30 minutes. Phenol extraction is repeated three to four times to execute deproteinization. Genome deposited through the ethanol precipitation is recovered, dissolved in 500 µl of sterile purified water containing 2.5 µg of ribonuclease, and left to stand at 42°C for 30 minutes. The phenol extraction is repeated two to three times to execute deproteinization. Genome deposited through the ethanol precipitation is recovered, and dissolved in 500 µl of TE. Thereafter, a positive control probe can be produced by measuring the concentration with an absorbance meter, and subjecting to digoxigenin labelling. Moreover, the positive control probe which may preferably used is one which permits to ascertain the hybrid formation when the positive control probe is subjected to dot hybridization on a membrane with 100 ng of U937 genome spotted thereon. When a negative control probe is required, it can be produced by any known method.

Further, the present invention also includes a kit for detecting and/or identifying causative microorganisms of infectious diseases by taking phagocytes from the clinical specimens containing active phagocytes, immobilizing the phagocytes so taken, treating the phagocytes to improve cell membrane permeabilities thereof, further treating the phagocytes to bare DNA in the causative microorganisms which might be existed in the phagocytes, *in situ* hybridizing DNA so bared with detective DNA probe(s) which can hybridize with such bared DNA under stringent conditions, and detecting and/or identifying the causative microorganisms based on signals so

detected, the kit comprises at least one or more enzyme selected from the group consisting of lysostafin, lysozyme, N-acetylmuramidase and zymolase used in the exposing step of the DNA, probe solution containing surfactant, and one or more DNA probe for detection. The kit includes, reagent for separating blood, enzyme pretreatment reagent, enzyme reagent, acetylation reagent, probe solution, blocking reagent, labelled antibody, labelled antibody diluent, coloring pretreatment liquid-1, coloring pretreatment liquid-2, coloring reagent, counter staining solution, PBS stock solution, hybridization stock solution, labelled antibody washing solution, coloring reagent washing solution, APS coated slide glass, probe dilution solution, buffer A and the like as demonstrated in the following Examples. Among these, it is preferred that at least the enzyme reagent and the probe solution are included. In addition, various reagents used in the present invention may be included for example, chloroform, ethanol, acetic anhydride, DMSO, PMSF, formamide, acetic acid, hydrochloric acid, sodium hydroxide and the like. Moreover, instrument and machine such as low speed centrifuge, incubator, counting chamber, shaker, humid box, incubator, light microscope, variable pipette, blood collection tube, tip, pipette, staining bottle, measuring cylinder, glass syringe, 0.2  $\mu$ m syringe top filter may be included.

Furthermore, the present invention provides a process for monitoring the gene of a foreign microorganism digested by a phagocyte included in a clinical specimen which contains a phagocyte derived from a living body. Moreover, the present invention provides a process for identifying the gene of a microorganism which becomes a candidate of the causative microorganism which a causative microorganism of sepsis or a causative microorganism of bacteremia is specified on the basis of the results identified.

It was revealed that when this process was applied in practice to diagnoses for blood of a variety of patients suspected as suffering from sepsis, causative microorganism

could be detected with about four times higher sensitivity compared to the blood culture process with no influence of the administered antimicrobial agent, and the identity of the detected microorganism strain was favorable. Furthermore, in comparison with the blood culture which requires three days or longer and approximately 14 days for the examination, an accurate result can be achieved by a simple operation within a short time period, i.e., about 8 hours, until the completion of the entire operation, according to the process of the present invention, an useful marker can therefore be realized in monitoring and the like in prognosis or diagnosis of an infectious disease such as sepsis, bacteremia or the like in which a rapid and favorable care is particularly necessary.

#### EXAMPLES

The present invention will be described in detail along with the following Examples, but as a matter of course would not be limited to the Examples because of the following disclosures.

##### EXAMPLE 1

##### Blooding and Preparation of Blood-Specimens

Twelve blood specimens (Specimen A-L) were obtained as a clinical specimen from the patients who are under the suspicion about sepsis. 10ml of heparinized venous bloods were obtained from each patient and were mixed with the blood components separative reagent (adjusted with sterilized-purified water to be 25ml as their final volume containing 225mg of Sodium Chloride and 1.5 g of Dextran (Molecular Weight: 200,000-300,000)) in the ratio of 4:1. Leukocyte fractions (the upper layer) were then obtained by leaving them at 37°C for 30 minutes. Leukocytes were appeared by centrifuging the leukocyte fractions in 160×g, at 4 °C and for 10 minutes. Pelletized leukocytes so obtained were suspended with 1 ml of sterilized-purified water and the suspension were immediately put into an isotonic state by adding thereto excessive amounts of PBS (prepared by diluting twenty-fold with sterilized-purified water the PBS Raw Solution

which have been adjusted with sterilized-purified water to be 120ml as their final volume containing 18.24 g of Sodium Chloride, 6.012 g of Sodium Monohydrogen Phosphate Didecahydrate and 1.123 g of Sodium Dihydrogen Phosphate Dihydrate). Such suspension were then re centrifuged in  $160\times g$ , at  $4^{\circ}\text{C}$  and for 10 minutes.

## EXAMPLE 2

### Fixation of Leukocytes

APS coated slides were employed wherein they were made by coating 3-aminopropyltriethoxysilane (APS, SIGMA) onto the slides (JAPAN AR BROWN CO., LTD., PRODUCTS ID. S311BL). Namely, such APS coated slides were produced by putting the slides (PRODUCTS ID. S311BL) into holders, immersing them into a diluted neutral detergent to wash the same, removing the detergent well therefrom with tap water then with purified water, leaving them under the higher temperature ( $100^{\circ}\text{C}$  or more) to dry the slides, and leaving the same under the room temperature to cool the slides. After then, these slides were immersed in Acetone containing 2% APS for one minute and were immediately rinsed gently with Acetone then with sterilized-purified water. These slides were then air-dried. Such performance was repeated and includes the immersion of these slides in Acetone containing 2% APS for one minute, immediate gentle rinse of the slides with Acetone and sterilized-purified water, and air-dry of the same. Finally, the slides were dried under the temperature of  $42^{\circ}\text{C}$  to realize the APS coated slides.

Pelletized leukocytes were prepared by centrifuging leukocyte fractions in  $160\times g$ , at  $4^{\circ}\text{C}$  and for 10 minutes. Then, leukocyte population was determined by adding small amount of PBS to the pelletized leukocytes, suspending the same, and counting the population with a hemacytometer. Leukocytes were duly mounted on the APS coated slides by smearing  $5\mu\text{l}$  of the leukocyte suspension into each well of the slides adjusted with PBS to be cell population of  $1\times 10^5$  cells/well, realizing the extended mono-layer of the leukocytes, and completely air-drying

the same. After then, they were immersed in Carnoy s fixative solution (prepared by mixing Ethanol, Chloroform and Acetic Acid in the volume ratio of 6:3:1) for 20 minutes and were immersed in 75% Ethanol solution for five minutes. Finally, they were completely air-dried.

### EXAMPLE 3

#### Treatment to Improve Permeability of Leukocyte Cell Membrane

They were immersed in PBS for 10 minutes, then in the solution prepared by diluting ten-fold a pretreatment reagent (prepared by mixing 1.25 g of Saponin, 1.25 ml t-octylphenoxy-polyethoxyethanol (specific gravity of 1.068-1.075 (20/4°C), pH (5w/v%) 5.5-7.5) and 25ml PBS Raw Solution, and adjusting with sterilized-purified water to be 50ml as their final volume), and were applied to a centrifuge for 10 minutes to accelerate thier permeability.

### EXAMPLE 4

#### Treatment of Bacterial Cell Wall with Lytic Enzymes

In order to bare DNA in the causative microorganisms, an enzyme solution was prepared by adding, per single slide, 1 ml of a reagent solvent (prepared by diluting 100-fold, with PBS, Dimethyl Sulfoxide (DMSO) containing 0.1 mol/l Phenylmethyl Sulfonylfluoride (PMSF)) to an enzyme reagent (N-acetyl-muramidase 1,000 Units/ml, Lysozyme 100,000 Units/ml and/or Lysostaphin 100 Units/ml), then 1 ml of which were dropped under the temperature of 37°C-42°C in the wet chamber onto the area where the leukocytes were smeared and were left for 30 minutes. Then, they were immersed in PBS containing 0.2 mol/l Hydrochloric Acid (prepared by adding hydrochloric acid to PBS Raw Solution, diluting 20-fold the same with sterilized-purified water and adujusting final concentration of Hydrochloric Acid to 0.2 mol/l), and were applied to a centrifuge for 10 minutes to accelerate thier permeability.



**EXAMPLE 5****Acetylation of Cell Membrane Proteins**

Slides were then immersed in the acetylation reagent prepared by adding Acetic Anhydride to the raw acetylation reagent (adjusted with quantum sufficient of sterilized-purified water to be 50ml as their total volume containing 7.46g of Triethanolamine and quantum sufficient of Hydrochloric Acid), diluting 10-fold with sterilized-purified water, and adjusting final concentration of Acetic Anhydride to 0.8%, and were applied to a centrifuge for 10 minutes. After then, they were completely air-dried by immersing them successively into 75%, 85% and 98% Ethanol for three minutes respectively.

**EXAMPLE 6****Alkalization of Bacterial DNA****(Denaturation of Double-Stranded DNA into Single-Stranded DNA)**

Slides were then immersed for three minutes in PBS containing 70mmol/l Sodium Hydroxide (prepared by adding Sodium Hydroxide to PBS Raw Solution, diluting 20-fold with sterilized-purified water, and adjusting final concentration of Sodium Hydroxide to 70 mmol/l). After then, they were completely air-dried by immersing them successively into 75%, 85% and 98% Ethanol for three minutes respectively.

**EXAMPLE 7****Hybridization**

Probe Solution (1.0 ng/ $\mu$ l) containing 15ng of DNA probes labelled with digoxigenin prepared with the diluted probe solution (containing 0.25% SDS, 600 $\mu$ l of Salmon Sperm DNA, 50 $\mu$ l of 100 $\times$ Denhart's Solution, 500 $\mu$ l of Hybridization Raw Solution, 2250 $\mu$ l of Formamide and 1000 $\mu$ l of 50% Dextran Sulfate) was applied to the smeared area and was left for two hours under the temperature of 37 $^{\circ}$ C-42 $^{\circ}$ C in the wet chamber. Probe Solution without SDS was employed as a control. DNA probes labelled with digoxigenin were prepared through nick translation methodology. Then, three stained bottles containing the hybridization detergent (prepared by mixing

Hybridization Raw Solution (prepared with sterilized-purified water to be 75ml as their total volume containing 13.15g of Sodium Chloride, 6.615g of Trisodium Citrate Dihydrate) in the ratio of Hybridization Raw Solution : sterilized-purified water : Formamide = 5:45:50) were provided and they were successively immersed therein at 42 °C for 10 minutes respectively.

After then, they were centrifuged for 10 minutes during which they were being immersed in PBS. As DNA probes labelled with digoxigenin, probes from *Staphylococcus aureus* of SA-24 (SEQ ID NO: 1), SA-36 (SEQ ID NO: 2) and SA-77 (SEQ ID NO: 3) as well as those from *Staphylococcus epidermidis* of SE-22 (SEQ ID NO: 4), SE-3 (SEQ ID NO: 5) and SE-32 (SEQ ID NO: 6) (See, Japanese Patent No.2798499) were employed. Also, as a probe from *Pseudomonas aeruginosa*, that of P2-2 (SEQ ID NO: 7) (See, Japanese Patent No.2965544) was employed. Then, as probes from *Enterococcus faecalis*, those of EF-1 (SEQ ID NO: 8), EF-27 (SEQ ID NO: 9) and EF-7 (SEQ ID NO: 10) (See, Japanese Patent No.2965543) were employed. Then, probes from *Escherichia coli* of EC-24 (SEQ ID NO: 11), EC-34 (SEQ ID NO: 12) and EC-39 (SEQ ID NO: 13), that from *Enterobacter cloacae* of ET-49 (SEQ ID NO: 14), and that from *Klebsiella pneumoniae* of KI-50 (SEQ ID NO: 15) (See, Japanese Patent No. 3026789) were also employed. Further, as probes from *Candida albicans*, those of CA-26 (SEQ ID NO: 16), CA-26-1 (SEQ ID NO: 17), CA-26-2 (SEQ ID NO: 18) and CA-26-3 (SEQ ID NO: 19) (See, Japanese Patent No. 2558420) were employed. The present probes were produced with the probes listed above through nick-translation methodologies.

#### EXAMPLE 8

##### Blocking

After *in situ* hybridization, blocking operation was performed. 1 ml of Blocking reagent (prepared with sterilized-purified water to be 10 ml as their final volume containing 2ml of Rabbit Normal Serum and 0.5 ml of PBS Raw Solution) per single slide was dropped onto the smeared area thereof and the slides

were left for 30 minutes. Then, Blocking reagent was removed.

**EXAMPLE 9****Reaction with Labelled Antibody**

Labelled antibody solution was prepared by diluting 50-fold the labelled antibody (prepared with 12.6  $\mu$ l of Buffer A (prepared with quantum sufficient of sterilized-purified water to be 100 ml as their total volume containing 746 mg of Triethanolamine, 17.5 mg of Sodium Chloride, 20.3 mg of Magnesium Chloride Hexahydrate, 1.36 mg of Zinc Chloride, 1000 mg of Bovine Serum Albumin and quantum sufficient of Hydrochloric Acid) to be 14  $\mu$ l as their total volume containing 1.05 Unit of alkaline-phosphatase-labelled anti-digoxigenin antibody solution) with the antibody dilution (prepared with quantum sufficient of sterilized-purified water to be 0.7 ml as their total volume containing 8.48 mg of Tris-(Hydroxymethyl)-Aminomethane, 6.14 mg of Sodium Chloride, and quantum sufficient of Hydrochloric Acid), then 10  $\mu$ l of the labelled antibody solution were dropped onto each of the smeared area, and they were left for 30 minutes. After then, they were immersed in the ten-fold diluted solution of the labelled antibody detergent solution (prepared with sterilized-purified water to be 100ml as their total volume containing 1ml of Polysolvate 20 and 50ml of PBS Raw Solution) and were centrifuged for 10 minutes during which they were being immersed in the detergent solution. This operation was repeated twice, then the samples were immersed in the treatment solution prepared by mixing Preliminary Treatment Solution 1 (prepared with quantum sufficient of sterilized-purified water to be 50ml as their total volume containing 6.06 g of Tris-(Hydroxymethyl)-Aminomethane, 2.92 g of Sodium Chloride, and quantum sufficient of Hydrochloric Acid) with equal amount of Preliminary Treatment Solution 2 (prepared with quantum sufficient of sterilized-purified water to be 50ml as their total volume containing 5.08 g of Magnesium Chloride Hexahydrate) and diluting 5-fold with sterilized-purified water. The samples were then centrifuged for 10 minutes during which they were being immersed in the treatment solution.

**EXAMPLE 10****Detection**

1 ml of chromogenic agent (Nitro Blue Tetrazolium (NBT)/ 5-Bromo-4-Chloro-3-Indolyl Phosphate (BCIP) Solution, pH 9.0-10.0: prepared with quantum sufficient of sterilized-purified water to be 10ml as their total volume containing 3.3mg of NBT, 1.65mg of BCIP, 99  $\mu$ g of N,N-dimethyl-formamide, 121mg of Tris-(Hydroxymethyl)-Aminomethane, quantum sufficient of Hydrochloric Acid, 58.4mg of Sodium Chloride, 101.6mg of Magnesium Chloride Hexahydrate) per single slide was dropped onto the smeared area of the slides by filtrating with a disposable syringe equipped with 0.2  $\mu$ m syringe top filter, and was kept in the dark and was left for 30 minutes at 37°C in a wet chamber. Then, they were immersed for five minutes in the solution prepared by diluting ten-fold the chromogenic agent cleaner (prepared with quantum sufficient of sterilized-purified water to be 50ml as their total volume containing 606mg of Tris-(Hydroxymethyl)-Aminomethane, quantum sufficient of Hydrochloric Acid and 186mg of Disodium Ethylenediaminetetraacetic acid Dihydrate) and were air-dried. Further, they were immersed in the solution prepared by diluting 10-fold the counterstain solution (prepared with quantum sufficient of sterilized-purified water to be 50ml as their total volume containing 50mg of Fast Green FCF (Food Dye Green No.3)) and in the 1% acetic acid solution. Thereafter, they were immersed again in the solution prepared by diluting 10-fold the cleaner aforementioned to remove excessive counterstain solution and were completely air-dried.

**EXAMPLE 11****Criterion**

When at least one expressed violet signal(s) is/are confirmed by the optical microscopy ( $\times 1,000$ ) on the subjected cells which were stained with the counterstain solution in single well, the sample was designated as positive. As the result of the present method, of twelve specimens, bacteria had been detected in the five specimens of Specimen A-SA (*Staphylococcus*

aureus), Specimen F and G-SE (*Staphylococcus epidermidis*), Specimen J-SE and EF (*Enterococcus faecalis*), and Specimen L-SA and CA (*Candida albicans*). In the meantime, when the same specimens were subjected to the hemoculture methodologies according to the known procedures, although the identical result was confirmed in Specimen A as a detection of SA, no bacteria had been detected in any of Specimen F, G, J and L. Accordingly, it became apparent that the present method could detect those rapidly with better sensitivity in comparison with the hemoculture methodologies.

With regard to the result on Specimen A-SA, the effect offered by adding SDS to the probe dilution is shown in Figure 1. Obviously from Figure 1, detection sensitivity on the signal could be improved remarkably with addition of 0.25% SDS. Turning to the other specimens, signals could similarly be detected clearly by adding SDS thereto. Meanwhile, the probes employed in this Example were those produced by performing nick translation on a combination of the base sequences from SA-24 (SEQ ID NO:1), SA-36 (SEQ ID NO:2) and SA-77 (SEQ ID NO:3).

## EXAMPLE 12

### Investigation on Optimum Cell Population of Leukocytes to be Smearred and Mounted

Optimum Cell Population of Leukocytes to be smeared onto the well (the circular well of 5mm diameter) of APS coated slides had been investigated. 10ml of heparinized healthy human bloods were collected, then, the leukocytes were obtained therefrom according to the procedures of Example 1. Leukocytes so obtained were suspended in the quantum sufficient of PBS and the leukocyte population per 1ml was determined with a hematimeter. A serial dilution commencing at (a)  $1 \times 10^8$  cells/ml and followed by (b)  $5 \times 10^7$  cells/ml, (c)  $1 \times 10^7$  cells/ml, (d)  $5 \times 10^6$  cells/ml, (e)  $1 \times 10^6$  cells/ml, (f)  $5 \times 10^5$  cells/ml and (g)  $1 \times 10^5$  cells/ml was prepared and 5  $\mu$ l of each population were smeared onto the slides.

Smeared slides were air-dried and were fixed with Carnoy's

fixative solution (See, Example 2). Immediately thereafter, they were stained with the counterstain solution aforementioned and were subjected to the criterion according to the method of Example 11. As a result thereof, the cell population of  $1 \times 10^8$  cells/ml was too many to detect and was not appropriate. Then, the cell populations of  $5 \times 10^6$  cells/ml or less were not appropriate, because little population were observed in the well. Accordingly, phagocyte density (X cells/ml) to be immobilized should be adjusted to the range of about  $5 \times 10^6$  cells/ml  $< X$  cells/ml  $<$  about  $1 \times 10^8$  cells/ml, preferably, to that of about  $1 \times 10^7$  cells/ml  $\leq X$  cells/ml  $\leq$  about  $5 \times 10^7$  cells/ml. It was concurrently apparent therefrom that leukocyte population (Y cells/well (5mm diameter)) to be immobilized in the single well of APS coated slides should be adjusted to the range of about  $2.5 \times 10^4$  cells/well  $< Y$  cells/well (5mm diameter)  $<$  about  $5 \times 10^5$  cells/well, preferably, about  $5 \times 10^4$  cells/well  $\leq Y$  cells/well (5mm diameter)  $\leq$  about  $2.5 \times 10^5$  cells/well. Experimental results on the samples (a)-(f) were illustrated in Figure 2 (a)-(f) respectively.

### EXAMPLE 13

#### Investigation of Lytic Enzyme to be Employed

Enzymatic conditions to lyse *Staphylococcus aureus* (ATCC 12600), *Staphylococcus epidermidis* (ATCC 14990), *Pseudomonas aeruginosa* (ATCC 10145), *Enterococcus faecalis* (ATCC 19433) and *Escherichia coli* (ATCC 11775) had been investigated. Lysostaphin (Bur. J. Biochem., 38, pp.293-300, 1973) was used as an lytic enzyme for *Staphylococcus aureus* and *Staphylococcus epidermidis*. As for *Enterococcus faecalis*, N-acetylmuramidase (Archs. Oral Biol., 23, pp.543-549, 1978), Lysozyme (SEIKAGAKU CORPORATION) was used. Then, as for *Pseudomonas aeruginosa* and *Escherichia coli*, PBS containing 70 mmol/l sodium hydroxide was used. All of these bacteria were inoculated into 5ml of BHI (Brain Heart Infusion) liquid medium (DIFCO) and were cultivated under 37 °C for eight hours or more. Cultivated solution were collected by centrifuging them in  $2,000 \times g$ , at 4 °C, for 10 minutes. Samples to be subjected were prepared by suspending

the collected bacteria in PBS.

Lysis activities were determined with a microplate reader by evaluating decreased density at 600nm absorbance on the subjected sample solution. As a result thereof, *Staphylococcus aureus* and *Staphylococcus epidermidis* were lysed with Lysostaphin. As for *Pseudomonas aeruginosa* and *Escherichia coli*, enzyme treatment was not necessary, because they were lysed with PBS containing 70 mmol/l sodium hydroxide. Then, as for *Enterococcus faecalis*, it was discovered that lysis activities therefor were improved by not relying on N-acetylmuramidase alone but on the combination of N-acetylmuramidase with Lysozyme. But, for example, when *Pseudomonas aeruginosa*, *Escherichia coli* or the like are the digested bacteria with phagocytosis, such enzyme treatments may not be necessary, because their cell walls are solved at the alkaline treatment, thereby, their genes are bared. Each enzyme to be employed in the pretreatment procedure of the present invention for solving the exogenous microorganism would be valid not only for the previously noted bacteria strain, but also for the other bacterial strains including those belonged to *Staphylococcus* genus, *Streptococcus* genus, *Bacillus* genus and *Micrococcus* genus. Such enzymes can be used as an individual enzyme, but to employ them as the mixed enzymes is quite useful. These results were illustrated in Figure 3, in particular, as Figure 3 (a) *Staphylococcus aureus* and *Staphylococcus epidermidis*, (b) *Pseudomonas aeruginosa* and *Escherichia coli*, and (c) *Enterococcus faecalis*.

#### EXAMPLE 14

##### Investigation of Enzyme Solution

##### (Investigation of Optimum Concentration of DMSO)

Since protease contained in the enzyme reagent degrades a form of leukocyte, enzyme activity of DMSO which is solvent of PMSF to be used to keep the form of leukocyte had been investigated. *Enterococcus faecalis* was inoculated in 50ml of BHI liquid medium noted previously and was cultivated, at 37°C, for eight hours or more. Bacterial cells/ were collected by centrifuging the cultivation liquid, at 4°C, in 2,000×g,



for 10 minutes, then were suspended in PBS and were subjected to heat-treatment in an autoclave (120°C, 10 minutes). Next, these were centrifuged, at 4°C, in 2,000×g, for 10 minutes, then discarding the supernatant, and the precipitates were suspended in 1 ml of PBS and were lyophilized. The lyophilized samples were suspended in 5 mmol/l Tris-HCl (pH 6.0) containing 0-10% DMSO then 2 mmol/l Magnesium Chloride and were designated as a sample on N-acetylmuramidase. *Micrococcus luteus* (JCM1464) was inoculated in 5ml of BHI liquid medium (supra) and was cultivated, at 37°C, for eight hours or more. Bacterial cells/ were collected by centrifuging the cultivated bacterial solution, at 4°C, in 2,000×g, for 10 minutes. Then the supernatant was discarded, then, bacterial cells/ were collected by rinsing the bacterial pellets though suspension of them in PBS and recentrifuging the same, at 4°C, in 2,000×g, for 10 minutes. The bacteria so collected were suspended in PBS containing 0-10% DMSO and were designated as a sample on Lysozyme. On the other hand, samples on Lysostaphin were also prepared by culturing and collecting *Staphylococcus epidermidis* substantially along with the procedure for the samples on Lysozyme, and suspending the cultured bacteria with PBS containing 0%~10% DMSO.

Enzymatic activities were evaluated by determining, with a micro plate reader, decrease of absorbance at 600nm on the subjected samples. Correlation between DMSO and the enzymatic activities were experimented in this example under the enzymatic titer of (a) N-acetylmuramidase 300 Units/ml, (b) Lysozyme 10,000 Units/ml and (c) Lysostaphin 50 Units/ml. When each enzymatic activity was evaluated with decrease of the density of bacteria suspension (O.D.=600nm) per predetermined time, there is little correlation between DMSO and N-acetylmuramidase activity, while 5 % or more of DMSO lowered activities of Lysozyme and Lysostaphin. There was no decrease on the enzymatic activities in DMSO concentration of 2% or less. Accordingly, concentration of DMSO to solve PMSF is adjusted to less than 5%, preferably 2% or less, more preferably about 1%. Results

were shown in Figure 4 (a)-(c) and in the following Table 3.

**TABLE 3**  
Correlation between DMSO and Enzymatic Activity  
(Density Decrease in Bacteria Suspension)

Amounts of DMSO Added (%)	N-acetylmuramidase O.D./5 minutes	Lysozyme O.D./3 minutes	Lysostaphin O.D./10 minutes
0 (control)	79.3±4.8	0.689±0.028	0.168±0.017
0.1	75.0±3.2	0.678±0.026	0.164±0.009
1	75.8±2.8	0.660±0.026	0.160±0.008
2	75.8±2.5	0.653±0.024	0.145±0.009
5	76.3±4.9	0.566±0.017	0.124±0.006
10	73.8±3.5	0.464±0.016	0.094±0.006

#### Example 15

##### Examination on Enzymatic Lysis Solution

##### (Examination on Optimal Concentration of PMSF)

Because protease included in the enzyme reagent deteriorates the morphology of leukocytes, effects of PMSF (manufactured by PIERCE), which is added for the purpose of retaining the morphology of the leukocytes, on enzymatic activity were examined. PMSF was dissolved in 100  $\mu$ l of DMSO (manufactured by Wako Pure Chemical Industries, Ltd), and diluted to 10 ml with PBS such that the final concentration of PMSF becomes none (0 mmol/l) to 1 mmol/l. To this solution was added proteinase K (manufactured by Boeringer Mannheim) such that titer of the protease becomes 0.2 unit/ml. Heparinized healthy human blood in an amount of 5 ml was collected, and leukocytes were obtained according to the process described in Example 1. Next, the leukocytes were suspended in an appropriate amount of PBS, and the cell number was measured using a counting chamber. Cell number was adjusted to about  $5 \times 10^4$  cells/well to about  $2.5 \times 10^5$  cells/well, and 5  $\mu$ l therefrom was

smearred on the well of the APS coated slide glass. After air drying, fixation was executed according to the method of Carnoy fixation described in Example 2. Using this sample, tests were performed according to the process described in Examples 3 to 11. As a consequence of performing the tests at the concentration of PMSF of 1  $\mu\text{mol/l}$  to 1  $\text{mmol/l}$ , effects were found at the concentration of 10  $\mu\text{mol/l}$  or greater, while deterioration of morphology of the leukocytes was completely suppressed at the concentration of PMSF of 0.1  $\text{mmol/l}$  or greater. The results are shown in Fig. 5, for (a): protease 0.2 unit/ml alone, (b): 1  $\mu\text{mol/ml}$  PMSF added, (c): 10  $\mu\text{mol/ml}$  PMSF added, (d): 0.1  $\text{mmol/ml}$  PMSF added, and (e): 1  $\text{mmol/ml}$  PMSF added, respectively.

#### Example 16

##### Examination of Optimal Titer of Lytic Enzyme, Zymolase

Optimal titer of zymolase for exposing DNA was examined through lysis of *Candida albicans*. *Candida albicans* was inoculated in YPD medium, and cultured over day and night at 30°C. Then, two types of the solutions were prepared: a solution of *Candida albicans* as a substrate suspended in PBS (substrate 1); and a solution prepared by Carnoy's fixation, immersing in 70% ethanol, air drying and suspension in PBS (substrate 2). Upon the reaction, a mixture of zymolase/ PBS: 0.5ml, substrate: 1.5 ml, M/15 phosphate buffer: 2.5 ml and sterile purified water: 0.5 ml, adjusted to give the total volume of 5.0 ml was used.

Thereafter, the reaction was allowed at 37°C for 2 hours, and the  $\text{OD}_{800}$  was measured. Furthermore, the concentration of zymolase (Zymolyase-100T) for use was 0 mg/ml, 0.01 mg/ml, 0.025 mg/ml, 0.05 mg/ml, 0.1 mg/ml, 0.25 mg/ml, 0.5 mg/ml, 1 mg/ml, 2.5 mg/ml and 5 mg/ml. Consequently, each  $\text{OD}_{800}$  value when the substrate 1 was used was 0.533, 0.521, 0.553, 0.554, 0.548, 0.417, 0.394, 0.288, 0.163 and 0.113, and each  $\text{OD}_{800}$  value when the substrate 2 was used was 0.445, 0.411, 0.359, 0.282, 0.232, 0.146, 0.115, 0.096, 0.08 and 0.057. It was proven that effectiveness was brought when both of the substrate 1 and substrate 2 were in the range of 0.5 mg/ml to 5 mg/ml, and particularly 1 mg/ml

to 5 mg/ml. That is, the amount of zymolase to be used is preferably 50 unit/ml to 500 unit/ml, particularly 100 unit/ml to 500 unit/ml.

### Example 17

#### Examination of Optimal Condition (Titer) of Enzymatic Treatment

##### (1) Production of Digested sample

##### [1] Preparation of U937 cell

U937 cells (monocyte established cell line: ATCC CRL-1593.2) were cultured in an RPMI 1640 medium (25 ml) within a cell culture flask (175 cm<sup>2</sup>) in a 5% carbon dioxide gas incubator at 37°C. Next, the U937 cell culture liquid was placed in a 50 ml centrifuge tube, and centrifuged at 4°C for 10 minutes at  $220 \times g$  to recover the U937 cells. Then, thus recovered U937 cells were suspended in 200  $\mu$ l of PBS, and the cell number was counted with a counting chamber. The cell number was adjusted to  $1 \times 10^4$  cells/ $\mu$ l to  $2 \times 10^4$  cells/ $\mu$ l.

##### [2] Preparation of Bacterial Digested sample

*Staphylococcus aureus* (ATCC 12600), *Staphylococcus epidermidis* (ATCC 14990), *Pseudomonas aeruginosa* (ATCC 10145), *Enterococcus faecalis* (ATCC 19433) and *Escherichia coli* (ATCC 11775) were inoculated in each 5 ml of BHI culture medium, and cultured at 37°C for 8 hours or longer. The cultured bacterial liquid was centrifuged at 4°C for 10 minutes at  $2,000 \times g$  to collect the bacteria. After discarding the supernatant, the bacterial pellet was suspended in 5 ml of PBS, and centrifugation was conducted once again at 4°C for 10 minutes at  $2,000 \times g$  to collect the bacteria. Thus collected bacteria were suspended in 5 ml of PBS and thereafter, 15 ml of bacterial liquids was produced prepared by diluting in PBS to give the turbidity (O.D. = 600 nm) of the bacterial liquid, which was measured with a absorbance meter, of 0.01 to 0.03 for *Staphylococcus aureus*, 0.01 to 0.03 for *Staphylococcus epidermidis*, 0.02 to 0.03 for *Pseudomonas aeruginosa*, 0.01 to 0.03 for *Enterococcus faecalis*, 0.02 to 0.03 for *Escherichia coli*, respectively. Thus produced

bacterial liquid was transferred into a separate 175 cm<sup>2</sup> flask for culture, and left to stand still at room temperature for 30 minutes. Fifty ml of heparinized healthy human blood was collected, and thereto was added the reagent for separating hemocyte at a ratio of 4 : 1, and left to stand still at 37°C for 30 minutes to yield the leukocyte fraction. Thus obtained leukocyte fraction was adjusted to 50 ml with PBS. The supernatant in the culture flask (*supra*) was gently discarded, and each 10 ml of the leukocyte fraction diluted in PBS was added to the flask followed by leaving to stand still at room temperature for 10 minutes. The supernatant in the flask for culture was discarded, and the leukocytes attached to the bottom of the flask were recovered in a 15 ml centrifuge tube with 10 ml of PBS containing 0.02% EDTA, and centrifuged at 4°C for 10 minutes at 140×g to 180×g to collect the leukocytes. Because contamination of erythrocytes was found in the collected leukocytes, precipitates of the leukocytes were gently suspended in 1 ml of sterile purified water to allow hemolysis, subjected to isotonization through adding 14 ml of PBS, followed by centrifugation once again at 4°C for 10 minutes at 140 × g to 180×g to collect the leukocytes. The collected leukocytes were suspended in PBS, and cell number was counted with a counting chamber to adjust to give 1×10<sup>4</sup> cells/μl to 5×10<sup>4</sup> cells/μl. These digested samples were referred to as SA digested sample, SE digested sample, PA digested sample, EF digested sample and EK digested sample.

### [3] Smear and Fixation

Each 5 μl of U937 cells prepared in Example 17 (1) [1] and each 5 μl of each bacterial digested sample produced in Example 17 (1) [2] were smeared on each well of the APS coated slide glass, and air dried. Next, after immersing the slide glass in the Carnoy's fixative described in Example 2 for 20 minutes, it was immersed in 75% ethanol for 5 minutes. After washing Carnoy's fixative and air drying, the slide glass was stored at 4°C until use in the test (see, Example 2). Next, pretreatment of the fixed sample was carried out according to Example 3.

(2) Standard and Process for Testing Digested sample[1] Cell Number

Cell number to be smeared and fixed on the slide glass of each bacterial digested sample was  $5.0 \times 10^4$  to  $2.5 \times 10^5$  cells/well, whilst cell number of U937 cells was  $5.0 \times 10^4$  to  $1.0 \times 10^5$  cells/well.

[2] Phagocytosis Rate

The bacterial digested sample smeared and fixed on the slide glass was stained with an acridine orange staining solution, and about 200 cells were randomly counted with a fluorescence microscope ( $\times 1,000$ ). Among the measured cells, cells including bacteria phagocytized within the cells (cells with any change characteristic in phagocytosis found in morphology, as shown by arrows in Fig. 6) were determined as positive cells, and the phagocytosis rate (%) was calculated according to the mathematical formula below.

$$\text{Phagocytosis rate (\%)} = [(\text{Positive cell number} / \text{Measured cell number}) \times 100]$$

Thus calculated phagocytosis rate (%) of each bacterial digested sample was 10% or greater.

[3] Test Process

The digested sample produced in Example 17 (2) [1] and [2] was employed as a specimen. The SA digested sample used had the phagocytosis rate of 23% with  $1.98 \times 10^5$  cells/well. The SE digested sample had the phagocytosis rate of 27% with  $1.74 \times 10^5$  cells/well. Moreover, the EF digested sample had the phagocytosis rate of 34% with  $6.40 \times 10^4$  cells/well.

Using the slide glass having each digested sample smeared thereon, the enzymatic pretreatment was performed according to the process described in Example 3. Next, the slide glass after completing the enzymatic pretreatment was placed in a humid box, and the reaction was allowed by dropping 1 ml of each enzyme solution prepared to give each titer on the smeared site of the specimen. Thereafter, the slide glass was immersed in PBS

containing 0.2 mol/l hydrochloric acid, and in 70% ethanol respectively, for 10 minutes, and air dried. After immersing this slide glass in PBS containing 70 mmol/l sodium hydroxide for 3 minutes, and in 70% ethanol for 10 minutes, it was air dried and stained with 1% acridine orange staining solution. Then, evaluation was made with a fluorescence microscope ( $\times 1,000$ ). For *Staphylococcus aureus* and *Staphylococcus epidermidis*, examination of the optimal titer was conducted with lysostafin. In order to examine the optimal titer when N-acetylmuramidase and lysozyme are used in combination for *Enterococcus faecalis*, examination on optimal titer of lysozyme was conducted in cases where N-acetylmuramidase was fixed at 100 unit/ml, and on optimal titer of N-acetylmuramidase in cases where lysozyme was fixed at 10,000 unit/ml. The determination was made as adequate when no bacterial body was identified in the leukocytes by the enzymatic treatment.

#### [4] Results

For the lysis of *Staphylococcus aureus*, as described in Table 4, sufficient effects were exerted at the titer of lysostafin of 1 unit/ml, however, upon lysis of *Staphylococcus epidermidis*, the titer of lysostafin of 10 unit/ml or greater was necessary. Therefore, the optimal titer of lysostafin was set to be 10 unit/ml to 100 unit/ml. In addition, for the lysis of *Enterococcus faecalis*, lysis did not occur with the titer of N-acetylmuramidase of 10 unit/ml or less when the titer of lysozyme was fixed at 10,000 unit/ml. In respect of lysozyme, when the titer of N-acetylmuramidase was fixed at 100 unit/ml, lysis did not occur with the titer of lysozyme of 1,000 unit/ml or less, as described in Table 5. Therefore, the optimal titer of N-acetylmuramidase was set to be 100 unit/ml to 1,000 unit/ml, whilst the optimal titer of lysozyme was set to be 10,000 unit/ml to 100,000 unit/ml. The results are shown in Fig. 7. In the Figure, depicted are states of: (a) the digested sample of *Staphylococcus aureus* prior to the enzymatic treatment, (b) the digested sample of *Enterococcus faecalis* prior to the enzymatic treatment, (c) the sample (a) following the enzymatic treatment,

and (d) the sample (b) following the enzymatic treatment.

Table 4: Optimal Titer for Enzymatic Treatment  
of Lysostafin on *S. Aureus*, *S. epidermidis*

Lysostafin Titer (U/mL) Digested Samples		0	0.1	1	10	100	1,000
SA Digested Sample	once	inadequate	Inadequate	adequate	adequate	adequate	adequate
	twice	inadequate	inadequate	adequate	adequate	adequate	adequate
	thrice	inadequate	inadequate	adequate	adequate	adequate	adequate
SE Digested Sample	once	inadequate	inadequate	inadequate	adequate	adequate	adequate
	twice	inadequate	inadequate	inadequate	adequate	adequate	adequate
	thrice	inadequate	inadequate	inadequate	adequate	adequate	adequate



**Table 5: Optimal Titer of Enzymatic Treatment  
of N-acetylmuramidase and lysozyme on *E. faecalis***

N-acetyl muramidase titer (U/mL) Digested Sample		0	1	10	100	1,000	10,000
EF Digested Sample	once	Inadequate	inadequate	inadequate	adequate	adequate	adequate
	twice	Inadequate	inadequate	inadequate	adequate	adequate	adequate
	thrice	Inadequate	inadequate	inadequate	adequate	adequate	adequate
Lysozyme titer (U/mL) Digested Sample		0	10	100	1,000	10,000	100,000
EF Digested Sample	once	Inadequate	inadequate	inadequate	inadequate	adequate	adequate
	twice	Inadequate	inadequate	inadequate	inadequate	adequate	adequate
	thrice	Inadequate	inadequate	inadequate	inadequate	adequate	adequate

Applications of these results obtained using the digested samples to the present invention could result in similar results. Therefore, the optimal titer of each enzyme as described above in the identification of a causative microorganism of an infectious disease in the clinical specimen of the present invention was also set similarly.

#### **Example 18**

##### Examination on Optimal Condition of Enzymatic Treatment (Temperature)

Using a slide glass including each digested sample smeared thereon, examination was conducted according to the process described in example 17 (2) [3]. Time period of the enzymatic treatment in this test was 30 minutes, and the temperature for examination was 4°C, 25°C, 37°C, 42°C, and 60°C. Moreover,

titer of each enzyme was N-acetylmuramidase (100 unit/ml, manufactured by Seikagaku Corporation), lysozyme (10,000 unit/ml, manufactured by Seikagaku Corporation), lysostafin (10 unit/ml, manufactured by SIGMA).

Determination was conducted according to the process described in example 17 (2) [3]. As a consequence, for *Staphylococcus aureus*, no bacterial body was found in the leukocytes in the range of temperature of 4°C to 60°C. For *Staphylococcus epidermidis*, although bacterial bodies remained in the leukocytes at the temperature of 4°C and 25°C, no bacterial body was found at 37°C or higher. Further, for *Enterococcus faecalis*, although bacterial bodies remained at the temperature of treatment of 4°C, 25°C and 60°C, no bacterial body was found at 37°C and 42°C. Hence, the optimal temperature for the enzymatic treatment was set to be 37°C to 42°C. The results are shown in Table 6.

Table 6: Optimal Temperature for Treatment of Enzyme Reagent

Temperature for Treatment (°C)		4	25	37	42	60
Digested Samples						
SA Digested Sample	Once	adequate	adequate	adequate	adequate	adequate
	twice	adequate	adequate	adequate	adequate	adequate
	thrice	adequate	adequate	adequate	adequate	adequate
SE Digested Sample	once	inadequate	inadequate	adequate	adequate	adequate
	twice	inadequate	inadequate	adequate	adequate	adequate
	thrice	inadequate	inadequate	adequate	adequate	adequate
EF Digested Sample	once	inadequate	inadequate	adequate	adequate	inadequate
	twice	inadequate	inadequate	adequate	adequate	inadequate
	thrice	inadequate	inadequate	adequate	adequate	inadequate

Applications of these results obtained using the digested samples to the present invention could result in similar results. Therefore, the optimal temperature of the enzymatic treatment in the identification of a causative microorganism of an infectious disease in the clinical specimen of the present invention was also set similarly.

#### Example 19

##### Examination on Optimal Condition of Enzymatic Treatment (Time)

Digested samples produced according to the process described in Example 17 (1) [1] and [2] were used as specimens. Time period of the examination was 0 minute, 10 minutes, 20 minutes, 30 minutes, 60 minutes and 120 minutes. Phagocytosis rate of the used SA digested sample was 18% with  $7.80 \times 10^4$  cells/well. Phagocytosis rate of the used SE digested sample was 34% with  $1.10 \times 10^5$  cells/well. Further, phagocytosis rate of the EF digested sample was 28% with  $1.30 \times 10^5$  cells/well.

Using the slide glass including each digested sample smeared thereon, examination was conducted according to the process described in example 17 (2) [3]. Temperature for the enzymatic treatment in this test was 37°C, and titer of each enzyme was 100 unit/ml for N-acetylmuramidase, 10,000 unit/ml for lysozyme, 10 unit/ml for lysostafin. Determination was conducted according to the process described in example 17 (2) [3]. As a consequence, for all of *Staphylococcus aureus*, *Staphylococcus epidermidis* and *Enterococcus faecalis* digested samples, no bacterial body was found in the leukocytes with the time period of the enzymatic treatment of 20 minutes or longer (inadequate at 0 minute and 10 minutes). Therefore, the optimal time period of the enzymatic treatment is at least 15 minutes or longer, preferably 20 minutes or longer, and still preferably 30 minutes to 60 minutes. The results are shown in Table 7.

Table 7: Optimal Time Period of Treatment of Enzyme Reagent

Time of enzyme-treatment (minutes)		0	10	20	30	60	120
SA Digested Sample	once	inadequate	inadequate	adequate	adequate	adequate	adequate
	twice	inadequate	inadequate	adequate	adequate	adequate	adequate
	thrice	inadequate	inadequate	adequate	adequate	adequate	adequate
SE Digested Sample	once	inadequate	inadequate	adequate	adequate	adequate	adequate
	twice	inadequate	inadequate	adequate	adequate	adequate	adequate
	thrice	inadequate	inadequate	adequate	adequate	adequate	adequate
EF Digested Sample	once	inadequate	inadequate	adequate	adequate	adequate	adequate
	twice	inadequate	inadequate	adequate	adequate	adequate	adequate
	thrice	inadequate	inadequate	adequate	adequate	adequate	adequate

Applications of these results obtained using the digested samples to the present invention could result in similar results. Therefore, the optimal time period of the enzymatic treatment in the identification of a causative microorganism of an infectious disease in the clinical specimen of the present invention was also set similarly.

#### Example 20

##### Examination on Optimal Condition of Enzymatic Treatment (Time)

In *in situ* hybridization reaction according to the present invention, concentration of the probe is an important factor which affects the hybridizing velocity. When the probe concentration is too low, the reaction velocity may be lowered, leading to the possibility of unclear signal. Furthermore, use of an excess amount of probe may result in grounds for nonspecific reaction.

Thus, optimal concentration was examined in connection with various probe solutions. First, the digested samples produced according to the process described in Example 17(1) [1] and [2] were used as specimens. The phagocytosis rate of the used SA digested sample was 24% with  $1.48 \times 10^5$  cells/well. The phagocytosis rate of the SE digested sample was 28% with  $2.07 \times 10^5$  cells/well. The phagocytosis rate of the PA digested sample was 11% with  $1.59 \times 10^5$  cells/well. In addition, the phagocytosis rate of the EF digested sample was 24% with  $1.72 \times 10^5$  cells/well. The phagocytosis rate of the EK digested sample was 12% with  $1.63 \times 10^5$  cells/well. Using the slide glass including each digested sample smeared thereon, examination was conducted according to the process described in Example 17(2) [3]. The probes for use were labelled with digoxigenin, and the concentration of each probe for *Staphylococcus aureus*, *Staphylococcus epidermidis*, *Enterococcus faecalis*, *Pseudomonas aeruginosa* and *Escherichia coli* was adjusted to 0.06 ng/ $\mu$ l, 0.6 ng/ $\mu$ l, 1.2 ng/ $\mu$ l, 1.8 ng/ $\mu$ l, 2.4 ng/ $\mu$ l, 3 ng/ $\mu$ l, respectively. The probe solution prepared to each concentration described above was used on the slide glass including the digested sample smeared thereon (see, Fig. 8), and examined according to the process described in Examples 3-11.

Consequently, the signal became unclear at lower concentration (0.06 ng/ $\mu$ l), and on the other hand, increase in background was observed at higher concentration (2.4 ng/ $\mu$ l and 3 ng/ $\mu$ l). Therefore, the concentrations of probes of SA, SE, PA, EF and EK were determined to be 0.6 to 1.8 ng/ $\mu$ l, preferably 0.6 to 1.2 ng/ $\mu$ l. Moreover, since an inadequate result was yielded at 0.06 ng/ $\mu$ l, while an adequate result was yielded at 0.6 ng/ $\mu$ l, it is preferably determined to be 0.1 ng/ $\mu$ l or greater.

Furthermore, since an inadequate result was yielded at 2.4 ng/ $\mu$ l, and an adequate result was yielded at 1.8 ng/ $\mu$ l, it is preferably determined to be 2.2 ng/ $\mu$ l or less. The results are shown in Tables 8-12 below.

Table 8: SA probe

Digested sample	Probe concentration (ng/ $\mu$ L)					
	0.06	0.6	1.2	1.8	2.4	3
SA digested sample	—	+	+	+	+	+
SE digested sample	—	—	—	—	+	+
PA digested sample	—	—	—	—	+	+
EF digested sample	—	—	—	—	+	+
EK digested sample	—	—	—	—	+	+

Table 9: SE probe

Digested sample	Probe concentration (ng/ $\mu$ L)					
	0.06	0.6	1.2	1.8	2.4	3
SA digested sample	—	—	—	—	—	+
SE digested sample	—	+	+	+	+	+
PA digested sample	—	—	—	—	—	+
EF digested sample	—	—	—	—	—	+
EK digested sample	—	—	—	—	—	+

Table 10: PA probe

Digested sample	Probe concentration (ng/ $\mu$ L)					
	0.06	0.6	1.2	1.8	2.4	3
SA digested sample	—	—	—	—	—	—
SE digested sample	—	—	—	—	+	+
PA digested sample	—	+	+	+	+	+
EF digested sample	—	—	—	—	—	+
EK digested sample	—	—	—	—	—	+

Table 11: EF probe

Digested sample	Probe concentration (ng/ $\mu$ L)					
	0.06	0.6	1.2	1.8	2.4	3
SA digested sample	—	—	—	—	—	+
SE digested sample	—	—	—	—	+	+
PA digested sample	—	—	—	—	+	+
EF digested sample	—	+	+	+	+	+
EK digested sample	—	—	—	—	—	—

Table 12: EK probe

Digested sample	Probe concentration (ng/ $\mu$ L)					
	0.06	0.6	1.2	1.8	2.4	3
SA digested sample	—	—	—	—	+	+
SE digested sample	—	—	—	—	+	+
PA digested sample	—	—	—	—	+	+
EF digested sample	—	—	—	—	+	+
EK digested sample	—	+	+	+	+	+

Applications of these results obtained using the digested samples to the present invention could result in similar results. Therefore, the optimal concentration of each probe described above in the identification of a causative microorganism of an infectious disease in the clinical specimen of the present invention was also set similarly.

### Example 21

#### Examination on Hybridization Temperature

Reaction temperature in the hybridization reaction is a parameter which affects the hybridizing velocity and stability of the hybrid. Because high temperature of the hybridization reaction is known to deteriorate the cell morphology, examination of the optimal temperature (4° C, 25° C, 37° C, 42° C,

50°C and 60°C) was performed.

First, the digested samples produced according to the process described in Example 17(1) [1] and [2] were used as specimens. The phagocytosis rate of the used SA digested sample was 31% with  $1.38 \times 10^5$  cells/well. The phagocytosis rate of the SE digested sample was 42% with  $1.95 \times 10^5$  cells/well. The phagocytosis rate of the PA digested sample was 14% with  $1.27 \times 10^5$  cells/well. In addition, the phagocytosis rate of the EF digested sample was 48% with  $1.05 \times 10^5$  cells/well. The phagocytosis rate of the EK digested sample was 17% with  $1.85 \times 10^5$  cells/well.

Using the slide glass including the digested sample and U937 cells smeared and fixed thereon (see, Fig. 9), examination was conducted according to the process described in Examples 3-11. Consequently, no stable signal was observed for each type of probe at the hybridization temperature of 4°C or less owing to the lowered hybridization velocity. Further, at 60°C, changes in cell morphology were detected, and thus no stable signal was observed. In addition, at 25°C and 50°C, detection could be executed better compared to at the temperature of 37°C and 42°C, although the signal was unclear. Hence, optimal temperature of the hybridization may be 25°C to 50°C, more preferably 37 to 42°C. The results are shown in Tables 13-17 below.

Table 13: SA probe

Digested sample	Hybridization temperature (°C)					
	4	25	37	42	50	60
SA digested sample	—	+	+	+	+	+
SE digested sample	—	—	—	—	—	—
PA digested sample	—	—	—	—	—	—
EF digested sample	—	—	—	—	—	—
EK digested sample	—	—	—	—	—	—



Table 14: SE probe

Digested sample	Hybridization temperature (°C)					
	4	25	37	42	50	60
SA digested sample	—	—	—	—	—	—
SE digested sample	+	+	+	+	+	—
PA digested sample	—	—	—	—	—	—
EF digested sample	—	—	—	—	—	—
EK digested sample	—	—	—	—	—	—

Table 15: PA probe

Digested sample	Hybridization temperature (°C)					
	4	25	37	42	50	60
SA digested sample	—	—	—	—	—	—
SE digested sample	—	—	—	—	—	—
PA digested sample	—	+	+	+	+	—
EF digested sample	—	—	—	—	—	—
EK digested sample	—	—	—	—	—	—

Table 16: EF probe

Digested sample	Hybridization temperature (°C)					
	4	25	37	42	50	60
SA digested sample	—	—	—	—	—	—
SE digested sample	—	—	—	—	—	—
PA digested sample	—	—	—	—	—	—
EF digested sample	+	+	+	+	+	—
EK digested sample	—	—	—	—	—	—

Table 17: EK probe

Digested sample	Hybridization temperature (°C)					
	4	25	37	42	50	60
SA digested sample	--	--	--	--	--	--
SE digested sample	--	--	--	--	--	--
PA digested sample	--	--	--	--	--	--
EF digested sample	--	--	--	--	--	--
EK digested sample	--	+	+	+	+	--

Applications of these results obtained using the digested samples to the present invention could result in similar results. Therefore, the optimal temperature of hybridization in the identification of a causative microorganism of an infectious disease in the clinical specimen of the present invention was also set similarly.

#### Example 22

##### Examination on Hybridization Time Period

The digested samples produced according to the process described in Example 17(1) [1] and [2] were used as specimens, and examination was conducted on the time period of hybridization of 10 minutes, 60 minutes, 90 minutes, 120 minutes, 180 minutes and 900 minutes. The phagocytosis rate of the used SA digested sample was 47% with  $1.45 \times 10^5$  cells/well. The phagocytosis rate of the SE digested sample was 47% with  $1.33 \times 10^5$  cells/well. The phagocytosis rate of the PA digested sample was 15% with  $1.91 \times 10^5$  cells/well. In addition, the phagocytosis rate of the EF digested sample was 41% with  $1.45 \times 10^5$  cells/well. The phagocytosis rate of the EK digested sample was 20% with  $1.23 \times 10^5$  cells/well.

Using the slide glass including the digested sample and U937 cells smeared and fixed thereon (same as one shown in Fig. 9), examination was conducted according to the process described in Examples 3-11. Consequently, although no signal was observed

with the time period of hybridization of 10 minutes, a signal was observed at 60 minutes or greater, and a stable signal was observed at 90 minutes or greater. Further, no alteration in detection of the signal was observed also with the time period of hybridization of 900 minutes. Therefore, it is preferred that the time period is at least 30 minutes or greater, preferably 60 minutes or greater, and more preferably 90 minutes or greater. More preferred optimal time period of hybridization may be set to be 120 minutes to 900 minutes. The results are shown in Tables 18-22 below.

Table 18: SA probe

Digested sample	Hybridization time (minutes)					
	10	60	90	120	180	900
SA digested sample	—	+	+	+	+	+
SE digested sample	—	—	—	—	—	—
PA digested sample	—	—	—	—	—	—
EF digested sample	—	—	—	—	—	—
EK digested sample	—	—	—	—	—	—

Table 19: SE probe

Digested sample	Hybridization time (minutes)					
	10	60	90	120	180	900
SA digested sample	—	—	—	—	—	—
SE digested sample	+	+	+	+	+	+
PA digested sample	—	—	—	—	—	—
EF digested sample	—	—	—	—	—	—
EK digested sample	—	—	—	—	—	—

Table 20: SE probe

Digested sample	Hybridization time (minutes)					
	10	60	90	120	180	900
SA digested sample	—	—	—	—	—	—
SE digested sample	—	—	—	—	—	—
PA digested sample	—	+	+	+	+	+
EF digested sample	—	—	—	—	—	—
EK digested sample	—	—	—	—	—	—

Table 21: EF probe

Digested sample	Hybridization time (minutes)					
	10	60	90	120	180	900
SA digested sample	—	—	—	—	—	—
SE digested sample	—	—	—	—	—	—
PA digested sample	—	—	—	—	—	—
EF digested sample	+	+	+	+	+	+
EK digested sample	—	—	—	—	—	—

Table 22: EK probe

Digested sample	Hybridization time (minutes)					
	10	60	90	120	180	900
SA digested sample	—	—	—	—	—	—
SE digested sample	—	—	—	—	—	—
PA digested sample	—	—	—	—	—	—
EF digested sample	—	—	—	—	—	—
EK digested sample	—	+	+	+	+	+

Applications of these results obtained using the digested samples to the present invention could result in similar results.

Therefore, the optimal time period of hybridization in the identification of a causative microorganism of an infectious disease in the clinical specimen of the present invention was also set similarly.

### Example 23

#### Influence of Surfactant Added to Hybridization Solution

The digested samples produced according to the process described in Example 17(1) [1] and [2] were used as specimens. When any of various surfactants (SDS, lauryl sarcosine, saponin, BRIJ35, Tween 20, Triton X-100) was added to the probe dilution solution followed by hybridization carried out according to Example 7, the detection sensitivity was dramatically enhanced by adding 0.25% SDS. In addition, the detection sensitivity could be improved by lauryl sarcosine, BRIJ 35 or Tween 20. The results are shown in Table 23 below.

Table 23

Surfactant	Signal detection
None added	+
SDS	+++
Lauryl sarcosine	++
Saponin	+
BRIJ 35	++
Tween 20	++
Triton X-100	+

Furthermore, as a consequence of using SDS at various concentrations, it was revealed that preferable concentration was 1% or less, more preferably 0.1% to 0.5%, and still more preferably 0.25%.

Applications of these results obtained using the digested samples to the present invention could result in similar results.

Therefore, also in the present invention, it is preferred that a surfactant, particularly SDS, is added at the step of *in situ* hybridization.

#### Example 24

##### Examination on Chain Length of Probe Used in Hybridization

*Staphylococcus aureus* probe (SA-24 (SEQ ID NO: 1)) and *Pseudomonas aeruginosa* probe (P2-2 (SEQ ID NO: 7)) were labelled with digoxigenin.

First, 1 µg of purified each type of the DNA probe was prepared to give the total volume of 50 µg with 5 µl of 10× L.B. (0.5 mol/l Tris-hydrochloric acid (pH 7.5), 50 mmol/l magnesium chloride, 5 µl of 0.5 mg bovine serum albumin), 5 µl of 100 mmol/l dithiothreitol, each 1 nmol of dNTPs (A, G, C), 0.5 nmol of digoxigenin-dUTP (Dig-dUTP), each 0.5 nmol of dTTP, 3 µl of DNase (amount corresponding to 25 mU, 75 mU and 200 mU), 1 µl of 10 U/µl DNA polymerase and an appropriate amount of sterile purified water. Digoxigenin labelling was performed at 15°C for 2 hours. After the labelling, the mixture was boiled for 5 minutes to terminate the reaction. The reaction liquid after the termination was injected into a spin column (CENTRI-SEP COLUMUNS CS901, PRINCETON SEPARATIONS, INC.), and centrifuged at 25°C for 2 minutes (3,000×g) to remove free nucleotides. Then, concentration of the eluate was measured by an absorbance meter. Electrophoresis was performed on a 3% agarose gel to confirm the size.

Next, DNA in the agarose gel was transferred to a nitrocellulose membrane by Southern blotting method. Then, the membrane was immersed in 2% blocking reagent (manufactured by Roche) for 30 minutes, and thereafter, alkaline phosphatase labelled anti-digoxigenin antibody in an amount of 1/5,000 was added thereto and the immersion was allowed for 30 minutes. Next, the membrane was washed twice by shaking in 100 mmol/l Tris-hydrochloric acid (pH 7.5) and 150 mmol/l sodium chloride for 10 minutes. Thereafter, washing was executed by shaking in

100 mmol/l Tris-hydrochloric acid (pH9.5) and 150 mmol/l sodium chloride for 10 minutes. Then, color development was conducted by immersing in an NBT/BCIP solution.

Finally, the membrane was immersed in purified water to stop the color development, and dried. Consequently, as shown in Fig. 10 for (a) use of the SA probe and (b) use of the PA probe, respectively, it was indicated that in cases where cleavage was conducted using 25 mU of DNase (in Figure, lane 1) such that the chain length distributes the base length of predominantly about 350 to about 600, high labelling efficiency was achieved. When thus resulting probe for detection was used in the process for detecting a causative microorganism of an infectious disease according to the present invention in which a digested sample or a clinical specimen from a patient suffering from an infectious disease was used to carry out hybridization, a signal could be detected with excellent sensitivity. Therefore, it was revealed that chain length of the probe used in the hybridization may be the base length of about 350 to about 600, and preferably the base length of about 350 to about 550.

#### Example 25

##### Examination on Probe used in Hybridization

*Escherichia coli* digested samples produced according to the process described in Example 17(1) [1] and [2] were used as specimens to examine on the probes for detection.

Probes for detection were prepared through labelling with digoxigenin as described in Example 24 from EC-24 (SEQ ID NO: 11), EC-34 (SEQ ID NO: 12) and EC-39 (SEQ ID NO: 13) such that they have the base length of about 350 to about 600, and used alone or in combination of those three, respectively. From thus obtained results, it was evident that the signal could be detected more clearly resulting in elevated sensitivity for (d) the mixed probe MIX prepared by mixing the three ((a) EC-24, (b) EC-34 and (c) EC-39), than for each (a) EC-24, (b) EC-34 or (c) EC-39 used alone, as shown in Fig. 11.

INDUSTRIAL APPLICABILITY

Since the *in situ* hybridization according to the present method can offer the stable signals within two hours or less, detection results can be presented very rapidly. Obviously, such rapid detection demonstrates the value of the present method in the rapid diagnosis of sepsis.



## CLAIMS

1. A method for detecting and/or identifying causative microorganisms of infectious diseases by taking phagocytes from the clinical specimens containing active phagocytes, immobilizing the phagocytes so taken, treating the phagocytes to improve cell membrane permeabilities thereof, further treating the phagocytes to bare DNA in the causative microorganisms which might be existed in the phagocytes, *in situ* hybridizing DNA so bared with detective DNA probe(s) which can hybridize with such bared DNA under stringent conditions, and detecting and/or identifying the causative microorganisms based on signals so detected, the method comprises at least one condition(s) to be selected from the following conditions (1)-(8) of;

(1) Cell density ( $X$  cells/ml) of the phagocytes to be immobilized is  $5 \times 10^6$  cells/ml  $< X$  cells/ml  $< 1 \times 10^8$  cells/ml,

(2) Lysostaphin is applied into the step to bare DNA in the titer of from 1 Unit/ml to 1,000 Units/ml,

(3) Lysozyme is applied into the step to bare DNA in the titer of from 1,000 Units/ml to 1,000,000 Units/ml,

(4) N-acetylmuramidase is applied into the step to bare DNA in the titer of from 10 Units/ml to 10,000 Units/ml,

(5) Zymolyase is applied into the step to bare DNA in the titer of from 50 Units/ml to 500 Units/ml,

(6) Surfactant is applied into the step of *in situ* hybridization,

(7) Such DNA probe(s) is/are one or more DNA probe(s) to be determined with their chain length of from 350 bases to 600 bases, and

(8) Concentration of such DNA probe(s) is from 0.1 ng/ $\mu$ l to 2.2 ng/ $\mu$ l.

2. The method according to Claim 1, wherein said step to bare DNA employs one or more enzyme(s) selected from Lysostaphin in the titer of 10-100 Units/ml, Lysozyme in the titer of 10,000-100,000 Units/ml, N-acetylmuramidase in the titer of 100-1,000 Units/ml and Zymolyase in the titer of 100-500 Units/ml.

3. The method according to Claim 1 or 2, wherein said step to bare DNA employs enzyme(s), and the enzyme(s) is/are subjected to a reaction to be performed under the temperature of 26°C-59°C for 15-120 minutes.

4. The method according to any of Claims 1-3, wherein said step to bare DNA further employs substance(s) to keep a form of the phagocytes.

5. The method according to Claim 4, wherein said substance to keep a form of the phagocytes is phenylmethanesulfonyl fluoride.

6. The method according to Claim 5, wherein concentration of said phenylmethanesulfonyl fluoride is  $10\mu\text{mol/l} \sim 10\text{mmol/l}$ .

7. The method according to any of Claims 4-6, wherein said substance to keep a form of the phagocytes is a substance dissolved into dimethylsulfoxide.

8. The method according to Claim 7, wherein said substance to keep a form of the phagocytes is a substance dissolved in dimethylsulfoxide, and concentration of the dimethylsulfoxide in a solution to be employed in said step to bare DNA is adjusted to less than 5%.

9. The method according to any of Claims 1-8, wherein said *in situ* hybridization step is performed by hybridizing DNA with DNA probe(s) under the presence of surfactant(s).

10. The method according to Claim 9, wherein said surfactant is an anionic surfactant.

11. The method according to Claim 10, wherein said anionic surfactant is sodium dodecyl sulfate.

12. The method according to any of Claims 1-11, wherein a hybridization reaction in said *in situ* hybridization step is performed under the temperature of 25°C-50°C for 30-900 minutes.

13. The method according to any of Claims 1-12, wherein said method comprises, prior to said immobilization step, a step to mount the phagocytes so taken onto the solid support, and the solid support is a slide coated with 3-aminopropyl triethoxysilane.

14. The method according to any of Claims 1-13, wherein the method employs, at said signal detection, a pigment to distinguish contrast between signals and cells/.

15. The method according to any of Claims 1-14, wherein said clinical specimen is blood.

16. A kit for detecting and/or identifying causative microorganisms of infectious diseases by taking phagocytes from the clinical specimens containing active phagocytes, immobilizing the phagocytes so taken, treating the phagocytes to improve cell membrane permeabilities thereof, further treating the phagocytes to bare DNA in the causative microorganisms which might be existed in the phagocytes, *in situ* hybridizing DNA so bared with detective DNA probe(s) which can hybridize with such bared DNA under stringent conditions, and detecting and/or identifying the causative microorganisms based on signals so detected, the kit comprises the following elements (1)-(2) of;

(1) at least one enzyme(s) to be employed in said step to bare DNA which is/are selected from Lysostaphin, Lysozyme, N-acetylmuramidase and Zymolyase; and

(2) at least one detective DNA probe(s).

17. A method for monitoring a gene of exogenous microorganisms digested with the phagocytes in the clinical specimens containing active phagocytes comprising the step of detecting the gene with *in situ* hybridization method employed in the method according to any of Claims 1-15, wherein the gene of exogenous microorganisms in the clinical specimens is monitored.

18. A method for diagnosing sepsis or bacteriemia comprising the step of identifying a gene of candidate causative microorganisms with *in situ* hybridization method employed in the method according to any of Claims 1-15, wherein the causative microorganisms for sepsis or bacteriemia are determined based on the identification results.

## S E Q U E N C E   L I S T I N G

&lt;110&gt; FUSO PHARMACEUTICAL INDUSTRIES, LTD.

TSUNEYA OHNO

<120> Improved Method for Detecting and Identifying  
Causal Microorganisms of Infectious Diseases

&lt;130&gt; 02P327W0

&lt;150&gt; JP 2001-165929

&lt;151&gt; 2001-05-31

&lt;160&gt; 19

&lt;210&gt; 1

&lt;211&gt; 10207

&lt;212&gt; DNA

&lt;213&gt; Staphylococcus aureus

&lt;220&gt;

&lt;223&gt; Designated as SA-24

&lt;400&gt; 1

aagcttatgg acctatttta ggtatattga ttagttggct tggattaatt tctggaacat	60
ttacagtcta tttgatctgt aaacgattgg tgaacactga gaggatgcag cgaattaaac	120
aacgtactgc tgttcaacgc ttgattagtt ttattgatcg ccaaggatta atcccattgt	180
ttattttact ttgttttcct ttacgcaa atacattaat aaattttgta gcgagtctat	240
ctcatattag acctaaatat tatttcattg ttttggcatc atcaaagtta gtttcaacaa	300
ttattttagg ttatttaggt aaggaaatta ctacaatttt aacgcacctt ttaagagggg	360
tattaatggt agttgtgttg gttgtatttt ggattgttgg aaaaaagtta gaacagcatt	420
ttatgggacg gaaaaaggag tgacatcgtg aaaaaagttg taaaatatat gatttcattg	480
atacttgcta ttatcattgt actgttcgta caaacttttg taatagtttg tcatgtcatt	540
ccgaataatg atatgtcacc aacccttaac aaagggacgt gttattgtaa ataaaattaa	600
agttacattt aatcaattga ataatggtga tatcattaca tataggcgtg gtaacgagat	660
atatactagt cgaattattg ccaaacctgg tcaatcaatg gcgtttcgtc agggacaatt	720
ataccgtgat gaccgaccgg ttgacgcac ttatgccaag aacagaaaaa ttaaagattt	780

tagtttgcgc	aattttaaag	aattagatgg	agatattata	ccgcctaaca	attttgttgt	840
gctaaatgat	catgataaca	atcagcatga	ttctagacaa	tttggtttta	ttgataaaaa	900
ggatattatt	ggtaatatata	gtttgagata	ttatcctttt	tcaaaatgga	cgattcagtt	960
caaactctta	aaagaggtgt	caaaattgaa	aaaagaatta	ttggaatgga	ttatttcaat	1020
tgcagtcgct	tttgtcattt	tatttatagt	aggtaaattt	attgttacac	catatacaat	1080
taaaggtgaa	tcaatggatc	caactttgaa	agatggcgag	cgagtagctg	taaacattat	1140
tggatataaa	acaggtgggt	tggaaaaagg	taatgtagtt	gtcttccatg	caaacaaaaa	1200
tgatgactat	gttaaacgtg	tcatcggtgt	tcctgggtgat	aaagtagaat	ataaaaaatga	1260
tacattatat	gtcaatggta	aaaaacaaga	tgaaccatat	ttaaactata	atttaaaaca	1320
taaacaagggt	gattacatta	ctgggacttt	ccaagttaaa	gatttaccga	atgcgaatcc	1380
taaatcaaata	gtcattccaa	aaggtaaata	tttagttcct	ggagataatc	gtgaagtaag	1440
taaagatagc	cgtgcgtttg	gcctcattga	tgaagaccaa	attgtttggt	aagtttcatt	1500
tagattcttg	ccatttagtg	aattttaaca	taatttcaat	cctgaaaata	ctaaaaatta	1560
atatgaaaca	aatacaacat	cgtttgtcgg	ttttaatact	gataaacgat	gttttatttt	1620
gttagtacca	caataaaagc	taagttcgaa	atgaacttat	aataaatcaa	tcacaatcac	1680
tttgtgttaa	aatatgtgtc	aaaggaagtg	agggtttgtc	atgacattac	atgcttattt	1740
aggtagagcg	ggaacaggta	agtctacgaa	aatgttgacc	gaaataaaac	aaaaaatgaa	1800
agcagatccg	cttgagatc	caatcatttt	aattgcgcca	actcaaagta	catttcaatt	1860
agaacaagcc	tttgtcaatg	atccggaatt	aaatggtagt	ttaagaacag	aagtgttgca	1920
ttttgaacga	ttaagtcatc	gtattttcca	agaagttggt	agttatagcg	aacaaaagtt	1980
atctaaagct	gcaacggaaa	tgatgattta	taacattggt	caagaacaac	aaaagtattt	2040
aaaactttat	caatcacaa	caaaatatta	tgggtttagt	gaaaaattaa	cagaacaaat	2100
tcaagatttt	aaaaaatatg	cagtaacgcc	tgaacattta	gaacacttta	ttgctgataa	2160
aaatatgcaa	actcgaacta	aaaataagtt	agaggatatt	gctttaatat	accgtgagtt	2220
cgaacaacgc	attcaaaaacg	agtttattac	tgggtgaggat	tcattacaat	attttattga	2280
ttgtatgccg	aatcagaggt	ggctaaaacg	tgctgatata	tatattgatg	gttttcacaa	2340
cttttcaacg	attgagtatt	taataatcaa	aggattaatt	aaatatgcga	gagtgtcaca	2400
atttatattga	cgacagatgg	taaccacgat	caatttagtt	ttttagaaaa	ccatcggaag	2460
tgttacgcga	tattgaagaa	atagcaaagt	aactcaatat	ttctattgaa	cgtcaatatt	2520
tcaaccaatt	atatcgcttc	aataatcaag	atttaaagca	tcttgaacaa	gaatttgatg	2580
tacttcaaata	caatcgagtg	gcatgtcaag	gtcatatcaa	tatttttagaa	tctgcgacta	2640
tgagagagga	aataaatgaa	attgcgcgac	gtatcatcgt	tgatattcgt	gataagcaat	2700
tacgatataca	agatattgca	attttatatc	gtgacgagtc	ttatgcttat	ttatttgatt	2760
ccatattacc	gctttataat	attccttata	acattgatac	aaagcgttcg	atgacacatc	2820
atccggtcat	ggaaatgatt	cgttcattga	ttgaagttat	tcaatctaata	tggcaagtga	2880
atccaatgct	acgcttattg	aagactgatg	tgtaaacggc	atcatatcta	aaaagtgcac	2940
acttagttga	tttacttgaa	aattttgtac	ttgaacgtgg	tatatacggg	aaacgttggt	3000

tagatgatga	gctattttaat	gtcgaacatt	ttagcaaaat	ggggcgtaaa	gcgcataaac	3060
tgaccgaaga	tgaacgtaac	acatttgaac	aagtcgttaa	gttaaagaaa	gatgtcattg	3120
ataaaatttt	acattttgaa	aagcaaatgt	cacaagcgga	aactgtaaaa	gactttgcaa	3180
ctgcttttta	tgaagtatg	gaatatttcg	aactgccaaa	tcaattgatg	acagagcgag	3240
atgaacttga	tttaaatggt	aatcatgaaa	aggcggagga	aattgatcaa	atatggaatg	3300
gcttaattca	aatccttgac	gacttagttc	tagtatttgg	agatgaacca	atgtcgatgg	3360
aacgtttcct	agaagtattt	gatatgtggt	tagaacaatt	agaatttgtc	atgattccac	3420
aaacattaga	tcaagttagt	attggtacga	tggatttggc	taaagtcgac	aataagcaac	3480
atgtttactt	agttggaatg	aaogacggca	ccatgccaca	accagtaact	gcacaaagtt	3540
taattactga	tgaagaaaag	aaatattttg	aacaacaagc	aaatgtagag	ttgagtccta	3600
catcagatat	tttacagatg	gatgaagcat	ttgtttgcta	tggtgctatg	actagagcta	3660
agggagatgt	tacattttct	tacagtctaa	tgggatcaag	tggtgatgat	aaggagatca	3720
gccccatttt	aaatcaaatt	caatcattgt	tcaaccaatt	ggaaattact	aacattcctc	3780
aataccatga	agttaacca	ttgtcactaa	tgcaacatgc	taagcaaacc	aaaattacat	3840
tatttgaagc	attgcgtgct	tggtagatg	atgaaattgt	ggctgatagt	tggtagatg	3900
cttatcaagt	aattagagat	agcgatcatt	taaatcaagg	tttagattat	ttaatgtcag	3960
cattaacgtt	tgacaatgaa	actgtaaaat	taggtgaaac	gttgtctaaa	gatttatatg	4020
gtaaggaaat	caatgccagt	gtatctcggt	ttgaagggtta	tcaacaatgc	ccatttaaac	4080
actatgcttc	acatgggtctg	aaactaaatg	aacgaacgaa	atatgaactt	caaaaactttg	4140
atttaggtga	tattttccat	tccgttttaa	aatatatata	tgaacgtatt	aatggcgatt	4200
ttaaacaatt	agacctgaaa	aaaataagac	aattaacgaa	tgaagcattg	gaagaaattt	4260
tacctaaagt	tcagtttaat	ttattaaatt	cttcagctta	ctatcgttat	ttatcaagac	4320
gcattggcgc	tattgtagaa	acaacactaa	gcgcattaaa	atatcaaggc	acgtattcaa	4380
agtttatgcc	aaaacatttt	gagacaagtt	ttagaaggaa	accaagaacc	aatgttacga	4440
attaattgca	caaacattaa	cgacaactca	aggtattcca	attaatatta	gagggcaaat	4500
tgaccgtatc	gatacgtata	caaagaatga	tacaagtttt	gttaatatca	ttgactataa	4560
atcctotgaa	ggtagtgcca	cacttgattt	aacgaaagta	tattatggta	tgcaaatgca	4620
aatgatgaca	tacatggata	tcgtttttaca	aaataaacia	cgccttggat	taacagatat	4680
tgtgaaacca	ggtggattat	tatacttcca	tgtacatgaa	cctagaatta	aatttaaatc	4740
atgggtctgat	attgatgaag	ataaactaga	acaagattta	attaaaaagt	ttaagctgag	4800
tggtttagtg	aatgcagacc	aaactgttat	tgatgcattg	gatattcggt	tagaacctaa	4860
attcacttca	gatattgtac	cagttgggtt	gaataaagat	ggctctttga	gtaaacgagg	4920
cagccaagtg	gcagatgaag	caacaattta	taaattcatt	cagcataaca	aagagaattt	4980
tatagaaaca	gcttcaaata	ttatggatgg	acatactgaa	gtgcaccatt	aaagtacaaa	5040
caaaaattgc	catgtgcttt	ttgtagttat	caatcggtat	gtcatgtaga	tggcatgatt	5100
gatagtaagc	gatatcgaa	tgtagatgaa	acaataaata	caattgaagc	aattcaaaat	5160
attaacatta	atgatgaatt	tgggggtgag	taatagatga	caattccaga	gaaaccacaa	5220

```

ggcgtgattt ggactgacgc gcaatggcaa agtatttacg caactggaca agatgtactt 5280
gttgcagccg cggcagggtt aggtaaaaca gctgtactag ttgagcgtat tatccaaaag 5340
attttacgtg atggcattga tgtcgatcga cttttagtcg taacgtttac aaacttaagc 5400
gcacgtgaaa tgaagcatcg tgtagaccaa cgtattcaag aggcatcgat tgctgatcct 5460
gcaaatgcac acttgaaaaa ccaacgcac aaaattcatc aagcacaaat atctacactt 5520
catagttttt gcttgaaatt aattcaacag cattatgatg tattaatat tgacccgaac 5580
tttagaacia gcagtgaagc tgaaaatatt ttattattag aacaaacgat agatgagggtc 5640
atagaacaac attacgatat ccttgatcct gcttttattg aattaacaga acaattgtct 5700
tcagatagaa gtgatgatca gtttcgaatg attattaaac aattgtattt ctttagcggt 5760
gcaaatccaa atcctacaaa ttggttggat caattggtga caccatacga agaagaagca 5820
caacaagcgc aacttattca actactaaca gacttatcta aagtatttat cacagctgcc 5880
tatgatgctt taaataaggc gtatgatttg tttagtatga tggatggcgt cgataaacat 5940
ttagctgtta tagaagatga acgacgttta atggggcgtg ttttagaagg tggttttatt 6000
gatatacctt atttaactga tcacgaattt ggcgcgcgtt tgcctaattg aacagcgaaa 6060
attaaagaag caaatgaaat gatggtcgat gccttagaag atgctaaact tcagtataaa 6120
aaatataaat cattaattga taaagtgaat aatgattact tttcaagaga agctgatgat 6180
ttgaaagctg atatgcaaca attggcgcca cgagtaaagt accttgcgcg tattgtgaaa 6240
gatgttatgt cagaattcaa tcgaaaaaag cgtagcaaaa atattctgga tttttctgat 6300
tatgaacaat ttgcattaca aattttaact aatgaggatg gttcgccttc agaaattgcc 6360
gaatcatacc gtcaacactt tcaagaaata ttggtcgatg agtatcaaga tacgaaccgg 6420
gttcaagaga aaatactatc ttgcatcaaa acgggtgatg aacataatgg taatttattt 6480
atgggttgag atgtaagca atccatttat aaatttagac aagctgatcc aagtttattt 6540
attgaaaagt atcaacgctt tactatagat ggagatggca ctggacgtcg aattgatttg 6600
tcgcaaaact cgttctcga aaagaagtac tgtcaacgac taactatata tcaaacatat 6660
gatggatgaa caagtcggtg aagtaaaata tgatgaagcg gcacagttgt attatgggtgc 6720
accatatgat gaatcggacc atccagtaaa cttaaaagtg cttgttgaag cggatcaaga 6780
acatagtgat ttaactggta gtgaacaaga agcgcatttt atagtagaac aagttaaaga 6840
tatcttagaa catcaaaaag tttatgatat gaaaacagga agctatagaa gtgcgacata 6900
caaagatatc gttattctag aacgcagctt tggacaagct cgcaatttac aacaagcctt 6960
taaaaatgaa gatattccat tccatgtgaa tagtcgtgaa ggttactttg aacaaacaga 7020
agtccgctta gtattatcat ttttaagagc gatagataat ccattacaag atatttattt 7080
agttgggtta atgcgtccg ttatatatca gttcaaagaa gacgaattag ctcaaattag 7140
aatattgagt caaatgatga ctacttctat caatcgattg taaattacat taatgacgaa 7200
gcagcagatg ctattttagt tgataaatta aaaatgtttt tatcagatat tcaaagttac 7260
caacaatata gtaaagatca tccggtgtat cagttaattg ataaatttta taatgatcat 7320
tatgttattc aatactttag tggacttatt ggtggacgtg gacgacgtgc aaacctttat 7380
ggtttattta ataaagctat cgagtttgag aattcaagtt ttagaggttt atatcaattt 7440

```



attcgtttta	tcgatgaatt	gattgaaaga	ggcaaagatt	ttggtgagga	aaatgtagtt	7500
ggtccaaacg	ataatgttgt	tagaatgatg	acaattcata	gtagtaaagg	tctagagttt	7560
ccatttgtca	tttattcttg	attgtcaaaa	gattttaata	aacgtgattt	gaaacaacca	7620
gttattttta	atcagcaatt	tggtctcgga	atggattatt	ttgatgtgga	taaagaaatg	7680
gcatttccat	ctttagcttc	ggttgcatat	aaagctgttg	ccgaaaaaga	acttgtgtca	7740
gaagaaatgc	gattagtcta	tgtagcatta	acaagagcga	aagaacaact	ttatttaatt	7800
ggtagagtga	aaaattgata	aatcgttact	agaactagag	caattgtcta	ttcttggtga	7860
gcacattgct	gtcaatgaac	gattaaacttc	accaaataccg	ttccatotta	tttatagtat	7920
tttatctaaa	catcaatctg	cgtcaattcc	agatgattta	aaatttgaaa	aagatatagc	7980
acaagttgaa	gatagtagtc	gtccgaatgt	aaatatttca	attatatact	ttgaagatgt	8040
gtctacagaa	accatttttag	ataataatga	atatcgttcg	gttaatcaat	tagaaactat	8100
gcaaaatggt	aatgaggatg	ttaaagcaca	aattaaacac	caacttgatt	atcaatatcc	8160
atatgtaaat	gatactaaaa	agccatccaa	aacaatctgt	ttctgaattg	aaaaggcaat	8220
atgaaagaag	aaagtggcac	aagttacgaa	cgagtaagac	aatatcgtat	cggttttcaa	8280
cgtatgaacg	acctaaattt	ctaagtgaac	aaggtaaacg	aaaaagcgaa	ttgaaattgg	8340
tacgttaatg	catacagtga	tgcaacattt	accattcaaa	aaagaacgca	tatctgaagt	8400
tgagttacat	cagtatatcg	atggattaat	cgataaacat	attatogaag	cagatgcgaa	8460
aaaagatatc	cgtatggatg	aaataatgac	attatcaata	gtgagtatat	tcgattattg	8520
ctgaagcaga	gcaagtttat	cgtgaattac	cgtttgtagt	taaccaagca	ttagttgacc	8580
aattgccaca	aggagacgaa	gacgtctcaa	ttattcaagg	tatgattgac	ttaatctttg	8640
ttaaagatgg	tgtgcattat	tttgtagact	ataaaaccga	tgcatttaat	cgtgcgcgtg	8700
ggatgacaga	tgaagaaatt	ggtacacaat	taaaaaataa	atataagata	cagatgaaat	8760
attatcaaaa	tacgcttcaa	acgatactta	ataaagaagt	taaaggttat	ttatacttct	8820
tcaaatttgg	tacattgcaa	ctgtagtatt	ttgattttca	aaagaataaa	aaataatttc	8880
gattaagtgc	aaagtccctg	tagcagaatg	aacacaactc	attttcaaaa	ttgtcttact	8940
tattttatttg	ttatttgata	acgaaaaaag	ttataatgtg	aattaagata	aagatgagga	9000
gttgagaatg	aatgaaattc	ttatcattca	agtataatga	caaaacttca	tatggcggtta	9060
aagtaaaacg	cgaagatgct	gtatgggatt	taacacaagt	atttgctgac	tttgacgaag	9120
gagattttcca	toctaaaaca	ttgttagctg	gtttacaaca	aaatcatact	ttagattttc	9180
aagaacaagt	acgtaaagca	gttgtagcag	cagaagatag	cggcaaagct	gaagactata	9240
aaatttcatt	taatgacatt	gaattcttac	caccagtaac	acctccgaat	aatgtgattg	9300
cttttggtag	aaattacaaa	gatcatgcga	acgaattaaa	tcatgaagta	gaaaaattat	9360
atgtattttac	aaaagcagcg	tcatctttta	caggagataa	tgcaacaatt	ccaaatcata	9420
aagatattac	tgatcaatta	gattatgaag	gtgaattagg	tattgttatt	ggtaagtctg	9480
gtgaaaagat	tccaaaagca	ttagcttttag	attatgttta	cggctataca	attattaacg	9540
atatcactga	tcgcaaagca	caaagtgaac	aagatcaagc	atttttatca	aaaagtttaa	9600
ctggcggttg	cccaatgggt	ccttatatcg	ttactaaaga	cgaactacca	ttacctgaaa	9660

```

atgtaaatat tgttacaaaa gttaacaatg aaattagaca agatggtaac actggcgaaa 9720
tgattcttaa aattgatgaa ttaatagaag aaatttcaaa atatgttgca ctactaccgg 9780
gagattatta ttgcaactgg tacaccagct ggcgttggtg caggtatgca accacctaaa 9840
tttttacaac caggtgatga agttaaagtg actattgata atattggaac gctgacaact 9900
tatatcgcta aataattatc atttaaaaag ctaaccaggt ctttatatag attggttagt 9960
tttttcttgc ttttctaaaa aggtgttaaa gataaattat ttataatgtt accattttga 10020
gatgaaagtg aaatattgat attaagaagt agttgattat ttacagcag attcacaata 10080
ttctaataag ggcaatgcaa atgtcatgtt cttcctctca aatatagaag tgtggtagaa 10140
tatatatctg tgtataatca aatctagatt aaattacaag caagtgggta ttaatcccaa 10200
gaagctt 10207

```

<210> 2

<211> 2082

<212> DNA

<213> *Staphylococcus aureus*

<220>

<223> Designated as SA-36

<400> 2

```

aagctttcta atctatcggt aatgatttgc tttaaaattg ggtogaagtt aattgaaggt 60
gtgaagtgtg tatctgtatt aataaccatg tcattcattt gctgcttcac tttgttaaca 120
agtcttccgt catataaaaa taatggtacg acaatcaatt ttgataccg tttcgagatg 180
ctttctaaat catgtgtaaa actaatctct ccatatagcg ttctcgcata agtaggttta 240
ttaatctgca aatgttgagc gcatatttgt aaotcttcgt gtgccttagt aaaatttoca 300
ttaatattgc cgtgtgcaac aaccataact ccaacttggt gttcgtcacc tgctaattgcg 360
tcacaaatac gttgttcaat taatcgtctc attaaaggat gtgtgccaag tggctcgctt 420
acttctacct ttatgtctgg ataccgtcgt ttcatttcat gaacgatatt oggtatatoc 480
ttgagataat gcattgcact aaagattagc aatggtacaa ttttaaaatg gtcaacccca 540
ctttgaatca acgtcgtcat taocgtctct aaatcctgat gctcacittc taaaaacgca 600
atatcatagt gatgtatata atcttttact aattcagaaa taaatgcttc taacgcttga 660
ttctgtcgtc cgtgcctcat gccatgtgca acaatgatat tccattcac atttaccaac 720
cctttcacac gtattgtata ccaaatacatt ttgtttttgt gaaaagaatc acattataat 780
gtaaaatcag ggaattccct gatgcctgta gtcatgcata ttcocttatac attttccott 840
tttgttaaat caaaaaaagc gaccgatata tgaatcccta ctcaacattt atttgagcaa 900
gcatcaatat atcggctcgt tgtagtgtat attattatct taaaatgggtg gttggcctaa 960
tattgtttcg tcaaagcgtc cgggtatcaa tactttgcgc atgatcacac ctaaatcgcc 1020

```

```

atcatcattt tcatgttcgc tgtatatattc ataacctctt ttttcataaa ttttaagtaa 1080
ccaaggatgc aatcttgcag atgtacctaa agtaactgcc gctgacttta acgtatctcg 1140
caaaaatgct cttcaacata agtaagtaat tggctaccat agcctttccc ttcatactca 1200
ggatttgtcg caaaccacca gacaaaagga tagcccgaaa tacttttcac acttccccaa 1260
ggatatctaa cagtaatcgt agatataatt tcatcatcaa ttgtcatgac aaatgtagta 1320
tttttatcta tttttcttt aacagcatct aaattagcat taactgaagg ccaatcaata 1380
cctagttctc ttagaggcgt aaatgcttca tgcattgagtt gttgcaattt ttctgcatct 1440
tgttcaactg cgagtogaat catcgttttt gtcattattaa tccccactct tttttaaatg 1500
atttaacat attttatatt taaaataaat atccatcaaa gtgtatcaat aaatttatca 1560
catgtcagaa agtatgttc atctgaatac accaatactc tcatgaaact tattaataat 1620
tactctctca acgtaaaaaa accattcaaa ttcattgaatg gtttggaaga atgattcatt 1680
gttaacgat ttaatcacta catcttaatt attgttgctc taaacgatta cgcttaccat 1740
ttaagaaagc ataaacgaga cctacaaaaa taccgccacc gacaaagtta cctaagaaag 1800
caaaaacgat atttttttaa acatgtaacc atgaaactgc atcaaggtta aagaatacca 1860
tacctgcata tagacctgca ttgaacacaa cgtgctcata tcccatgtat acaaagacca 1920
cgacaccaca agctatgaag aatgcctttg ttaagccgcc tttgaattgc atagagatga 1980
aaataccaat attaataaag aagttacaga aaataccttt tgtaaaaata ttcaaccatg 2040
ttgaatcaac agtctttttc tgaactaaag ctgttaaago tt 2082

```

<210> 3

<211> 2885

<212> DNA

<213> *Staphylococcus aureus*

<220>

<223> Designated as SA-77

<400> 3

```

aagcttttga ttaatttggg ctttaaagta ttoccaatta taattcttca tgattttctt 60
attggatttc gaatttgggt tcatgcattg ttgcctcaaa gaacatgctg aacagtcac 120
gcattcatat agcttgaagt cactttttaa accatatcta tcattacggt atgcatatct 180
tttaaaacct attottttgt tattaggaca tataaattca tcattaagtt cgtcatattt 240
ccaattttga gtgttaaaaa tgtaactttt aaactttcta gttttatctt taataaacat 300
gccatacgtg ataagtggcg ttttattaaa acatctataa tagccatata gttttgctca 360
ctatcataac tgcattcagct acattaaactc tggtaatacc gaggatttga atcattgtta 420
aaaatggaat taaagttcta gtatctgttg ggggttgaaa taggtcatag gataaaaaa 480
ttgagaattt gtcgtatttt gtaaatgtg tctggotta agttggccat ttttcatatg 540

```

```

gtcttccttc atttcataa aagttgcac atgacagcc cagaaagcta tttctatctt 600
taagaatcca tttttgttct tcataattat tttttcttcc ggaataatca tcaaatttct 660
ttttgaactt cttaatctca gttctttttt acgggtctgt tttctaattt gagcactctt 720
cgttctaaat agaatgattt aaatcttcga tttcttttat ctaaatgact accaattaaa 780
tctattttct ctcgtgattt tgaatacttt tcttccacac aaatgtatat ctattggcat 840
tagctttctac ttatgtacca tcaataaaaa ttgaattatt atcaataaga ttttgcttta 900
aacattgact atggaactga ataaataaag attcaattaa cgcacagta ttaggattca 960
ctctaaaacg attaatagtt ttataagaag gtgtttgatc ttgagctaac cacatcatto 1020
gaatactgtc atgaagtaat ttctctattc tacgaccaga aaatacagat tgagtatatg 1080
catataagat gatttttaac atcatttttg gatgatagga tgttgcgcca cgatgatgtc 1140
tgaattcatc gaattcgcta tcaggatcg tttcaacaat ttcatttaca tatcgcgaaa 1200
tatcatttta aggaattcta acagaagttt ctattggtag tgtaagttgg gcaaagtgtc 1260
ttattttttt aaagtatgta aaagtaaaat tacatgttaa tacgtagtat taatggcgag 1320
actcctgagg gagcagtgc agtcgaagac cgaggctgag acggcaccct aggaaagcga 1380
agcattcaat acgaagtatt gtataaatag agaacagcag taagatattt tctaattgaa 1440
aattatctta ctgctgtttt tttagggatt tatgtcccag cctgttttat tttcgactag 1500
tttgggagaat ttattgacat tcacattatt taaacggcaa caaagattgt tttattttga 1560
taggcattat atggtgttaa aaaatttgca tgaaaattaa aaaatgcttc gttcaggaag 1620
gtgtcgtaat ttacctattt gotgaatgaa gcattttatt tttaaatatg atagccaata 1680
taacaagcta taaatccaat gatgaattgt aaaagtgaat aattgagaaa aagggttaata 1740
tcaaattttg gtgtcatcat taatgtaagt tccttggcta acgttgagaa agttgttaag 1800
ccacctaaaa aaaccggtga caaagaacgc agggaaacct gagattgaaa ttgataggcc 1860
tatagttaat ccaattaaaa aactaccaac tagatttact atcaatgttg cgataggtaa 1920
ctttgaagta aatttatgat taaaataatc agtaatggca cttctagcaa ttgcgcaaaa 1980
accgcgcca atcatgacta aaatgattga tatcatgata aaccaccacc tagttttata 2040
ccgacgtaac ataacaaaat accaaagaca taacttgta cagcatatag tagtaaagtt 2100
ataaattggt gatgatcaaa catatgtatt aattctaatt gaaatgttga aaaagtcggt 2160
aaagcaccaa gaaaaccagt cgtaatatgt ttttttaggg tcggatgggt tgaaaaaat 2220
gcaattgtta aggotgttag caatcccatt acaaaggcac cagtcaaatt ggctatcagt 2280
gttccgattg gaaaacctcc gtcagtatto agaaaagaaa tgaggtaacg taataaagcg 2340
cctaaagcac caccgataaa aatatataca tattgcattt ggttcacctc gaaaagaagt 2400
agtttgaatt taaaaaagag gttttggcaa caogacgaca aaaattgtog atgcattatc 2460
aaacctcatt atatgttata tcttgttgta taactatagc gattagatgc atagtatatga 2520
tttcgaaaat ctaatatattt ttatacgcaa caacgtoatc aaattgtttt actcattata 2580
gcatgataca ttgtattgtt ttgtattaac gctacattga cattttatct tttttaaata 2640
aaaccgaatg tacgacaatt gaaaagatat gtactaaaat aacaattaga ataatccaag 2700
gcaaactttt actogcaatt ctaatccaat ctgcacagc ctttagtgat ttaattgaac 2760

```

gatctgcaaa aattatagac aaaattagta caattgagtt aataacactg cagaaaagta 2820  
 ttaatttaaat aaaagaatta aaaaatccac ttaggaaaac gttatttgta ttaaagaaaa 2880  
 agctt 2885

<210> 4

<211> 8654

<212> DNA

<213> *Staphylococcus epidermidis*

<220>

<223> Designated as SE-22

<400> 4

aagcttgttt tattgcttag ttatatctcc aataacactc attttatatg tacgtattgc 60  
 caaaaaaaaaat tatctataca gtaataagta tgaaatgaga actggaataa tcattggtat 120  
 tattgcttta attctagtaa ttatgcaagg gtttcacttt aactgggcta ttattcctat 180  
 ttctatctat ggtcatcagt ttgtattttt cgctggaatt attttaagtc ttgttggtat 240  
 attcttttaa cgtatagaat ttgtaggagt tggttacta ttttgtcaaa aacatagatg 300  
 caatggtaac tgacccggaa attgcacagt tttctcttt agcaatttgg attatacttg 360  
 ttgtgctaatt cattttttat acgatacgtt tatctgaacg cactaaatca tcatcatata 420  
 caaagattta aactcagaaa atatgctaga catatctttc tgagtttttt aatttattaa 480  
 aatatatcat ttgtttacca tataagtttg ttttagaaaa tgaatcacta ttttaataata 540  
 caaataaattt aattacactg aaaataacct aaaagcgtaa cactatttta atatgggtat 600  
 ataaatgact aaagggaggt gccaatgata ataaaattca aatttgtaat cagattgaac 660  
 ttaactatat tgatgaaggc gaaggcatcc ccatcatttt aattcatgga ttagatggaa 720  
 acttggcagg atttaaagat ttaaaaaatg aactcaagaa gcagtataga gtaattactt 780  
 atgatgtcag aggtcatgga aaatcttcac gaacagaatc atatgaatta aaagatcatg 840  
 ttgaagattt aaatgattta atgggagcat taaatatoga ttctgcacat attttaggac 900  
 atgatatggg gggcatcatt gcgagtgaat ttactgaaaa atatcaatat aaagtgatta 960  
 cattgacaat tgtttcggcc aaaagtgaag acattgcaaa tggtttcaac aaattaatgg 1020  
 ttgattacca agaagaatta gcaggcttta ataaatctga ggcaatgatt attttattct 1080  
 ctaaattatt taaagagaaa gataaagcaa tgaaatgggt atcaaagcca aaaattatac 1140  
 aatagaccaa ctccggaaga aagtgcatt gcagtacgtg cattgcttaa tattaaagat 1200  
 ttaactcgtg ttcatacata tgtgtccata cctactttta ttgtgaatgg taagtatgac 1260  
 ccaactatac aaaataaaag tcattatgat atggatcaat attatgatca agttacaaaa 1320  
 attgtatttg ataattcagg acatgcacca catatcgagg aaccagaaaa attcctgaaa 1380  
 ctctacttag attttgtag ttaaaaaata agaacataaa taaaaacct taaatgatta 1440

ttgtcggaaa atcatttgag ggtttttag tagcagtaaa gtttggactc agatcactat	1500
cgtattaaact taataaaaga gtaaaacagt cttatctttc ataagtgaag gaaatatctg	1560
tttnactccc tagccattat acttcatttc attatttgc tctgtgatac ggttggtttac	1620
tcgtttaagt aaatcatcga tttttttacg ctgcttagaa tctactaaga ttaaaacagt	1680
tctttcatcg tgttcattac gttttttatt aaagtaattt tcttgagata aatttttaac	1740
agctttaaca acttgagggt gtttataatt taagtgattg ataatatctt taagataata	1800
ttcctctttt ttattctcac taatataagt taatactgca aattcttcaa agctgattga	1860
gaattctttt ttaattattc cttttaatct gtcagcataa gtgaccatag ctaataattc	1920
aaagcagtcg ttgatttttg aaatagccat taatgaaacc tccctattta tatcatatcc	1980
ataaatctta aaacccatct ttttaaatat aaagatagtt aattatatta ttgaattaag	2040
attacttgga tactataccc taatttatta atttatactt atttttctta tgaaaatacg	2100
aaagtgtccg tcataatata gtattaatat aaatttaaag aatatattta atgctatatt	2160
atttagttaa ttataactaa ataaaattaa gaagtaaaac aataagtgtt tataaaacaa	2220
attatctttt aaagtttata ctggaattag caatgtagca ttgctatat tcaaaaaaat	2280
aagattgttt ctaattttcc ttaatttaaa aaaaattata ctaaaaagaa tacttttttg	2340
aaagaatttt actaacattt tttatatata aatgtttatt aatttagaag taggattttt	2400
aacaactttt tcatctatca ataagccttt agttatatta atataccacc tttttaaact	2460
ctttttgtat gttacttctc tttttgtaga attaaaacat agcgtttttg aacaatagct	2520
gacgtaggta actctatgtc atttgaggct aatttgattt taaagtgtgt tccaatttga	2580
tgattgggtt gtgtagaaag taaaatgtcg taatatgaga cgccattttt tatttttgat	2640
ggtatattcg aaatttcttt aattttacta gtaaattgag tgttgtoact agatgttaca	2700
gaaatatttt gatttatttt taataaatc aactcagatt ctgatatatt agcacgaata	2760
atacgttcgt tgcattaat ttgcactatc ttttcgtttg gttttgaagg gatagaatta	2820
atatatgaaa tacttccatt aattggtgaa aataaagtgg atttaattga ggatttagtt	2880
tgaatcattt gtaatttttag ctgattaagg aatgaataat aatgtaaato atttttagaa	2940
tttaaagttt tgttgttacg ttcatcacta agtgtatttt ggagttcttc atataaatga	3000
tctttttcat aattgtaata ttctaacact ggagtgtttt tagatacttt gctatgattt	3060
tttactaaaa gtttttgagg ttgtcctaaa gtgggagtggt agtagaaaat atagctgtta	3120
agaggggctt gtataaccagt tgttgaaagg agtaatttgg gctttgcttt tatagttttt	3180
atatttttaa tatctctgtt tttagaagtt aatttagaga aagtaatgta actaaaacta	3240
caagttgtga gaatgaaaat gaatagtaat gaagaaataa cgatgcgttg cttgggtcatg	3300
gatgttcaac tcataatatt attgtgaggt tattatacac tattattttt aatgaaatat	3360
attaattttt aataagcatt acttttggtt tgtatattgt tttatttcaa aaaataaagt	3420
aatcaattt aataaattga aaaatagaag gctatcttta attttaaaat atatgattct	3480
acataaatgt tactataaga agaactactc ataaaaactg ccaacaaaga caaatcttt	3540
gttggcagtt ogaaatagac atttatttgt atgaggaato tacattaata taagcggata	3600
atttttattc agaataagga attttaaata atcgtaataa aataatacct atagctatac	3660

ataataatcc	acctaactta	cgtgatgtta	ttttgttttt	aggtgaaccc	aacaaaccga	3720
aatgatcgat	aataataccc	ataatcattt	ggcccatcat	agcaattata	gtagttaaag	3780
ctgctcctaa	gaaaggcatt	aaaataatat	tagatgttac	gaatgccatt	cctagtatcc	3840
ctccaataaa	ataaatagat	ttaatcttac	ctagtgtttt	atgagtagat	gatattttca	3900
gactacgatt	aaataactaat	gttaatatata	ataacgctat	tgtaccaacg	ctaaatgata	3960
tgagtgaagc	aaatatggat	gagtgtgtgt	gttgagccag	tgtgctgttg	attgttgttt	4020
ggattggcgg	acgaaaccaa	atacgaatcc	aataagcaac	cagaatacta	ttggtgtatt	4080
cttatgtota	ttaacaggat	gtctacgaac	ataattcata	aatataattc	cagtaattaa	4140
aaatataatt	ccaacacctt	taaataatgt	aaaagattgt	tgatgggcgc	ccaataatcc	4200
aaatgtatca	atgattacac	ccataataat	ttgccctgta	accgtaataa	caacagtaag	4260
tgctgcgcct	aatcttggtg	ataataataa	gtttccagtt	aaatagataa	cacctaatag	4320
tcctcctagg	acccaagtat	agttaagtgt	ttgcttagaa	aagaattctg	gtgttaatac	4380
ttgtggatga	ataatgatat	taagcacaag	taagcatatt	gttcgcacag	caaaagatat	4440
ggttgaagca	taaaaagatg	aacgggtaaa	ttggcttagc	cttgagttga	ttgaagtttg	4500
aataggaagt	aacatgccaa	caaaaattcc	taaaagatat	agaaaaaaca	atgataaaaa	4560
ccaactttct	caatttaata	tgattatcat	accattcata	atcatgtttc	taaaatgatt	4620
gagccataag	caaagtatag	aaataagttg	tgaatgttcc	gaggtgtcat	acagccgata	4680
ctattttgat	gaatcattat	aataaaatgc	acattaaaca	agtttttagaa	ttaaaaaaag	4740
cgagacatca	ttttgaattt	gatatctcac	ttcatattiaa	taaaagaaca	atgtaaatta	4800
agttcttttt	tagacttgaa	caatttttaa	aaatttgttc	ttcgataagt	cttttttatg	4860
atttttagtac	tttaaataaa	gcgtcaaaaa	taatgtttta	tgaattaatt	tttatottca	4920
aatataacag	ttgtcctttt	atcaataagt	tgtgcagcat	aaattttgac	aggctttccc	4980
aaactaaatc	ttaaaatgtc	taattctaaa	atgtctaatt	ctaaaagttg	gttcataactt	5040
totttaatta	attgtttctgt	agtaatagcg	ttaaaatcgg	gtaatagtaa	tttgacgggt	5100
ttattaagat	ttgatttaaa	tacgagttcc	aaagtttttg	acatactgat	gtatcctcct	5160
taaattaaag	attctgtttt	aacgatctcg	actttgtcat	actcttcgcc	actgaacggt	5220
caatgatgga	acgaaaagat	ttgatttgat	cattagaaac	aagcggatta	atgttagaaa	5280
aacgacgcct	atgttcgact	actttacctt	cagaattatg	tttgatttga	gtaaagataa	5340
togtcaacttg	attgacttca	ttcataataa	aaacctcttt	cactatataat	atcgaaatag	5400
attgaaaaaa	aaggacacat	tttttgaaaa	atataggcaa	atgcotttga	tgtgatacaa	5460
acgtcattta	tcattaatta	tgaaacctgt	tttagaaggt	atatgaggta	agtagaattg	5520
ttagtttgta	aaagaaaaaa	ttggaacctg	atattttaaa	taaccaactt	aaaagattga	5580
tcagtgtota	aaattactat	ttatatatga	attaaaatat	taagatctcc	caatatgaga	5640
atgaattagt	ttaagtttat	cgatgattga	aaaattatag	cctcatggat	tctatottat	5700
ataaaaataaa	gttctattcc	cttttgata	taaataagaa	tagttacctt	tttgtgatat	5760
gccaatccag	aaaaaaaagcg	acagtgcctg	aatctatgta	tgctcaataa	actcattcaa	5820
atcaactagc	aatatcaaat	cataaatcgt	gttgcaccat	aataaggatt	aaaacctgtt	5880

agtttaacta	atttaagaaa	aacatttgat	tatottctot	ttcaatcggg	aatattaatt	5940
tctatcattc	aacaatattt	tggatatcag	ataacttaag	aaatattgag	atttattgaa	6000
atacgatatg	tttcaaactg	ccatacaatg	attacactta	ataaatgatt	acacttaata	6060
taaatgtaaa	aagaaaagga	ggggttaa	gagtttagta	tatcttatgg	cgactaattt	6120
attagtcatg	ctcatagttt	tattcactct	gagtcacgt	caactaagaa	agggtgcggg	6180
ctatgttgca	ttaatagctc	ctattgtgac	atctacatat	tttattatga	aaataaccaga	6240
tgtgattcga	aataagttta	ttgctgttcg	attaccatgg	atgccttcaa	ttgatattaa	6300
tttagattta	agattagatg	gtttaagttt	aatgttcggc	ttaattattt	cgctaatagg	6360
tgtgggtgta	tttttttatg	ctacgcaata	tttatccac	agtaaggaca	atottcctag	6420
atttttcatt	tatttactat	tatttatgtt	cagtatgatt	ggcattgtaa	tagctaataa	6480
taccatctta	atgtatgtat	ttgggaact	cacaagtatt	tctcattct	tgcttatatc	6540
ctattggtag	aataatgggtg	aaagtcgaatt	aggcgccatt	caatctttca	tgattacagt	6600
gtttgggtggg	ctagcgttat	taacaggatt	tatcatttta	tatatcatta	caggaacaaa	6660
cacaattact	gatattctaa	tcaacgcaat	gcaatttcac	gacatccttt	atttatacca	6720
atgattttga	tgctattatt	agggtctttt	accaaactcg	cacaatttcc	gtttcatatt	6780
tggttaccaa	aggccatggc	agcacctaca	ccagtaagt	cttatcttca	ttcggaaca	6840
atggtaaagg	ctggaatctt	ttactatatt	agatttacac	ctttattggg	acttagtaat	6900
gtttatattt	atacagtac	atttgttggt	ctaataacta	tgttatttgg	atctttaact	6960
gctttacgac	aatacgaact	aaaagggtata	ctcgttatt	ctacaataag	tcaattaggt	7020
atgattatga	caatggtagg	tctaggtggc	ggttatgctc	agcacacatc	agatgaattg	7080
tctaagtttt	atatttttagt	tttatttgot	ggcttattcc	atttaatgaa	tcatgcgggt	7140
tttaaagtgtg	cattatttat	gggcgttggt	atcattgatc	acgagtcggg	aacacgtgat	7200
attcgtttgc	taaatgggtat	gcgtaaagtc	tcccctaaaa	tgcatattgt	catgttgctc	7260
gctgcattat	ctatggcagg	tgctcctttt	ttaaatgggt	ttttaagtaa	ggaaatgttt	7320
ttagattcgt	taactaaagc	aaacgaactt	gatcaatatg	gcttcgtatt	aacgtttgtg	7380
attatttcaa	taggtgtcat	cgcgagtata	ttgactttta	cttatgcaat	ttacatgata	7440
aaagaaacat	tctggggaaa	ttacaatata	gaaaaattta	aacgtaaaca	aatacatgaa	7500
ccatggcctat	ttagtttacc	agctgtgatt	ttaatgttac	tcattccagt	tatcttcttt	7560
gttccaaaacg	tttttgga	ctttgttatt	ttgcccga	ccagatctgt	atctgggata	7620
gggcggaggt	tgatgcattt	gtgccacata	tttctcagtg	gcatgggtgtg	aatctccatt	7680
aattttaaga	tagtgtatat	attggactat	tttagctota	gtgtgattgg	aaagagggtta	7740
cgcacaaat	aatcaaaagt	gctcgattac	agtggctatc	ggaaatttat	agagaatttg	7800
aattatactc	agcccggtgt	atacgtgcat	tgatgaataa	taaattgaat	tattacatca	7860
tgattacatt	atttattttt	gtagctattg	tagttatgga	tatttgactg	tgggttttcc	7920
tcatgtactc	agcttcatat	tagttctttc	ggaccgttgg	aagttatctt	atcagttgta	7980
acattgatta	tccgcatctt	attaatcttt	attcgtcaac	gactaacgat	ggtgggtattg	8040
aatggaatga	ttggattcgc	agttacatta	tattttattg	caatgaaagc	tccagattta	8100



```

gctttaacac agttagtgt tgaaactatt acgacaatct tatttattgt tagtttttcg 8160
agactaccta acatccctcg agttaaggca aatttaaaaa aagagacctt caaatcatt 8220
gtgtcacttg ttatggcatt gacggtggtg tcaacttattt ttgttgctca acaagcagat 8280
ggtatgcctt caattgctaa attttatgaa gatgcatatg aacttacagg tggaaaaaat 8340
attgtcaatg ctatactagg tgacttcaga gcttttagata ctatgtttga aggactagt 8400
ttaatcatag ctggattagg tatttatacg ttacttaatt acaaagatag gagggggcaa 8460
gatgaaagag aatgatgtag tacttaaatc agttacaaaa attgtagtgt ttattttgtt 8520
aacatttggg ttttatgtat tttttgctgg ccataataat ccagggtggtg gctttattgg 8580
tggttgatt tttagctcgg catttatctt aatgtttctt gcctttgatg taaatgaagt 8640
gttgaaaaaa gctt 8654

```

<210> 5

<211> 2362

<212> DNA

<213> *Staphylococcus epidermidis*

<220>

<223> Designated as SE-3

<400> 5

```

aagcttcaca acttgaaaat atagcacaaa cattaaagga tttaggtaga aaacgagcaa 60
ttttaattca tgggtgcaaat gggatggatg aggccacgct ttctggtgaa aatatcattt 120
atgaagttag cagcgaaaga gcattaaaaa aatatagttt aaaagcagaa gaagtcggtt 180
tagcttatgc aaataatgac acgttgatag gtgggttcacc tcaaacaaat aaacaaattg 240
cattgaatat cctaagtggc acggatcact caagtaaacy agatgtagtt ttgttaaatg 300
ctggaattgc tttatatgtt gctgagcaag tggaaagtat caaacatggc gtagagagag 360
cgaaatatct cattgataca ggtatggcaa tgaacaata tttaaaaatg ggaggttaag 420
taatgactat tttaaatgaa attattgagt ataaaaaac tttgcttgag cgtaaatact 480
atgataaaaa acttgaaatt ttacaagata acggaaatgt taagaggaga aagctgattg 540
attcacttta actatgatag aacattatca gttattgctg aaataaaatc gaaaagccca 600
tctgtacctc aattaccgca acgtgatctt gtccaacaag ttaaagatta tcaaaaatat 660
ggtgctaatt ctatttcaat attaaactgat gaaaaatact ttggcggtag ttttgaacga 720
ttaaatcagt tatcaaagat aacatcggtt coagttttat gtaaagattt tattattgat 780
aaaattcaaa tagatgttgc aaaacgagct ggtgcatota ttattttatt aatagtaaatt 840
attttaagtg atgaccaatt aaaagaattg tattcatatg caacaaacca taatttagaa 900
gctotagtag aagttcatac aattagagaa cttgaacgtg cacaccaaatt taacctaaa 960
attattgggtg ttaataatcg tgatttaaaa cgatttgaaa ccgatgttct acatacaaat 1020

```

```

aaattactta agtttaaaaa gtctaattgc tgctacattt cagagagtgg cattcataca 1080
aaagaagatg ttgagaaaat agtagattca agtattgacg gtttacttgt aggggaggca 1140
ttaatgaaaa caaatgactt aagtcagttt ttgcctagt ttaaagttaa agaagaatct 1200
ctatgatagt taaattttgt ggttttaaaa ccgaaagtga tattaagaaa attaaaaaat 1260
tagaagttga tgcagtaggg ttatacatt atccogatag taagagacat gtctcactga 1320
aacaattaaa atatttggct aaaatagtgc cagatcatat agagaaagta gtgtcgtagt 1380
aaatcctcaa atgtccacca taaagagaat aattaatcaa actgatatta acacaatcca 1440
attacatgga aatgaaagca ttcaattaat tagaaatatt aagaaactta attcaaaaat 1500
aagaatcata aaagcaattc cagcaacaag aaatttaaat aataacattc aaaagtataa 1560
agatgagata gactatgttt attatagata caccatcaat cacatacggg gggacaggtc 1620
aaagttttga ctggaaatta ttaaaaaaaa taaaggcgtt gattttctca ttgcggtggt 1680
ttggattttg aaaagataaa acgattagaa atatattcat ttggacaatg tggttatgac 1740
atctcaactg gcattgagtc acataatgaa aaagatttta ataagatgac tcgaatatta 1800
aaatttttga aaggagacga atgattaatg aaaattcaaa cagaagtaga tgaattgggc 1860
tttttcggtg aatatggtgg ccaatatgta cctgaaacat tgatgccagc tattattgaa 1920
cttaaaaaag catatgagga cgcgaaatca gatactcact tcaagaaaga atttaattat 1980
tatttaagtg aatatgttgg tagagaaacg cctttaacat ttgctgaatc atacacaaaa 2040
ttgttaggtg gtgccaaaat atatcttaaa agagaagact taaatcacac tgggtgctcat 2100
aaaattaata acgcgatagg acaggcacta ttagctaaaa ggatggggaa aactaaatta 2160
gtagccgaaa cagggtgctgg tcaacatggt gtagcaagtg ccaccatcgc tgctttattc 2220
gatatggatc ttattgtttt catgggaagt gaagatatca aacgtcaaca acttaacgta 2280
tttagaatgg aattgctagg agctaaagta gtgtctgtgt cagatgggca aggaacacta 2340
tcagatgctg taaataaagc tt 2362

```

<210> 6

<211> 5024

<212> DNA

<213> Staphylococcus epidermidis

<220>

<223> Designated as SE-32

<400> 6

```

aagctttttg atttttaaag aaaaaattaa acaagggggc attgcttatg gtcaatagaa 60
gaaagatatc aattattggc gcgggacata cagggtgggac tctagcattc attcttgac 120
aaaaggaatt aggagatatt gtgttgattg aacgccagca atcagagggt atggctaaag 180
gaaaggcgtt agatatttta gaaagcggac ccatttgggg gtttgacaca totgtacatg 240

```

gttcagtaaa	tatagaagat	attaaagatt	cagacatagt	ggtgatgact	gcaggtatac	300
ctaggaaatc	aggaatgaca	aggagaagaa	ttagttcaaa	ctaatagaaca	aatagtacga	360
gaaactgcat	tacaaattgc	aacgtatgca	cctcattcaa	taattattgt	attgactaat	420
ccggttgatg	ttatgacata	tactgcattt	aaagcatcag	gttttcctaa	agaacgtatt	480
attggtcaat	ctggaatttt	agacgctgca	agatatcgaa	cttttattgc	tcaagaactt	540
aacgtgtctg	tcaaagatgt	aaatgggttt	gttttaggtg	gacatggtga	tacgatgtta	600
cctttgatta	ataacacaca	cattaatggg	attccagtta	agcatcttat	ttctgaagaa	660
aagattgatc	aaattgttga	acgtacacgt	aagggtggtg	cagaaattgt	tgcattacta	720
ggtcaaggct	cagcatatta	tgcaccagca	actgctatat	atgaaactat	agatgcaatt	780
tttaatgatc	ggaaacggtt	attaccaagt	attgcttata	tagagggaga	atacggttgt	840
tcagatatatt	gtttcggagt	tcctactata	ataggatatc	aaggaataga	aaagattata	900
gaggtagata	tgaataatga	tgagtatcaa	caactacaac	actctgcgca	agatgtgagt	960
gaagtcaaaa	actcactaaa	attcaaataa	ataattatga	agttctacat	cttaaattgt	1020
tagatttttg	tgaaaattgt	gtaaagggtg	ttttttcggt	gatttataaa	agcgctttct	1080
tgatataatg	aacatatatt	catagaataa	ggagacgatt	aaaatggcta	aaggggacca	1140
atatcaagct	catactgaaa	aatatcatga	gtaaaaagtc	taaaaaaagt	tataaacctg	1200
tgtggattat	cattagtttt	attattttta	ttacaatctt	gttattaccc	acaccagcag	1260
gattacctgt	aatggctaaa	gcagcactag	ctattttagc	tttcgctgta	gttatgtggg	1320
ttacagaagc	agttacttat	ccagtttctg	caacattaat	tttaggatta	atgatacttt	1380
tactagggtt	aagtccagtt	caagatttat	ccgaaaaact	tggaaaccta	aaagtggcga	1440
cataatacta	aaaggtagcg	atatttttag	aacgaataac	gcgcttagtc	acgcttttag	1500
tggtttttca	acctcagccg	tagcacttgt	agctgcagca	ttattttttag	cagtagctat	1560
gcaggaaacc	aatttacata	aacgacttgc	attattttgt	ctatcaattg	ttggaaataa	1620
aactagaaat	atagtcattg	gtgctatttt	agtatctatt	gttctagcat	tctttgtacc	1680
atcagctaca	gcacgtgctg	gtgcagttgt	cccaatatta	ctgggaatga	ttgctgcatt	1740
taatgtgagt	aaggatagta	gacttgcttc	attattaatt	attactgctg	tacaagcagt	1800
ttcgatatgg	aatataggta	ttaaaaacgg	ctgcagcaca	aaatattgta	gccatcaatt	1860
ttattaacca	aaatttagga	catgatgtat	catggggaga	gtggttttta	tatctgcgcc	1920
gtgggtcaatc	attatgtcta	tagctcttta	ttttataatg	attaagttta	tgccacctga	1980
acatgatgca	attgaagggtg	gaaaagagtt	aattaaaaag	gaacttaata	aattaggacc	2040
agtoagtoat	agagaatggc	gactaattgt	gatttcagtg	cttttatatt	ctctggtcga	2100
ctgagaaagt	attgcatccg	attgatccag	cttcogattac	actagttgct	ctaggtatta	2160
tgctaatgcc	aaagattggg	gttattaact	ggaaagggtg	tgaaaagaag	attccttggg	2220
ggacgattat	agtattttgg	gtaggaatct	cacttggtta	tgtattaact	aaaacaggag	2280
ccgctcatgg	ttagtgatca	acatttggtt	gatgggtctt	aaacatttac	cgatcatagc	2340
aactattgcg	ttaattacct	tatttaatat	attaatacat	ttaggttttg	caagtgcac	2400
gagcttagcc	totgcgttaa	tacctgtgtt	tatttctttg	acttcaacgc	taaatttagg	2460

tgatcatgct attgggttttg tattaataca acaatttgtg attagttttg gtttcctact	2520
acotgtcagt gcaccacaaa atatgcttgc atatgggtact gggactttta ccgtaaagga	2580
ttttttaaag acaggtatac ctttaacgat agtaggttat attttagtta tcgtatttag	2640
tttaacgtat tggaaatggc ttggtttagt gtaagtaaaa gatttaggta ttaaaatgat	2700
aattataaat gtctcgtaaa gtttaatat ttaactttac gacacatttt ttataaactc	2760
gtggcaagtt aatcttaata gttgaaatgt atcgataaaa aaatatatga atgtaaatag	2820
aatttagtat tagagaataa caaaaaattg atgttaggtg gtaaaatcta atggctatag	2880
gtgtcatatt aaatagagtt tttaggctaa ataataatcc attatttgat tatatatata	2940
gtaataaaga atctataaat cattgttatt ttattattcc aactgaagag tttgaagaag	3000
aagcaaaaaa gaaagcacaa tactattatg ggtccataca gaagtttatg tatgaactac	3060
aacgatatga tatagaacct tttttgatgt cttatgataa attaatagac ttttgtaaaa	3120
aacaagctat agacaaagtt gttgttgcag gtgatattat gagttatcat cacgaagaat	3180
atgacatttt acatcaaagg aaacgattta aacaagctaa tattcaagta atatcattaa	3240
gagcaaatca ttatttttaac ccccgcaaaa cacataataa acaaggggaa ccatataaag	3300
tatttaccag tttttataga aaatggcgtc ottacttaat gattagagat gaatatgact	3360
atcatttaga agatatttca aaggittgtg tgaaatctca acataaaatt aaagaagatt	3420
atcattcata tgggtataagt gaacgtgatg ttcaaaatcg ttggtctgaa tttttatctc	3480
aagatatcga aaattataaa gaaaacaggg aatacttgcc tgaagtatta acaagccaac	3540
taagtattta cttagcttat ggaatgatag atattataca atgttttcaa cgatttactt	3600
caaaattatg ataaaaatga acaaaattac gaaactttta tacgtgaatt gatttttaga	3660
gagttttatt atgtattaat gaccaattat cccgaaacag ctcatgttgc ttttaaagaa	3720
aaataccaac aattgaaatg gtcttataat gaagagaatt ttaaaactgtg gaaagatggg	3780
aatactgggt ttccaattat tgatgcagca atggaggaac ttaaaacaac tggatttatg	3840
cataatcgca tgagaatggg agtttctcaa tttttaacta aagatttggt tattgactgg	3900
atttgggggtg agtcattttt caaaacaaaa ttaatagatt atgatgcagc ttcaaatggt	3960
cacggatggc agtggtcagc ttctactgga acagatgctg taccatactt tagaatgttt	4020
aatcctataa gacaaagcga gcgttttgat aataatgcac gatataataa aacttacatt	4080
ccaagattaa atcaggtaga tgctaagtat ttacacgata ctcataaatt cgagcaacaa	4140
ataaaggggc aagggtgtga aataggtaaa gactatccta aacaaatgat tgatcacaaa	4200
gaaagtagac aacgtgtaat gtcagaatto aaagctatag attaaataaa aaagatctga	4260
acaacatgat atagggtgtt agatctttat ctagttacat aaaaaagcaa acatgaatta	4320
aaatatattc taacaaagtt aaaatataca tatatttaag atttaattta gttttcaaag	4380
gtacttccca atttgtataa cggggctcat aataaaataa ttgcatcaaa tataatccta	4440
tcctaacgg taaacacatt aataaaatag ctttagtata actccatcct atttgatgcc	4500
ataaatgacc tatcataagt tgaataatga tgagacatac cattaaaatt acttcaatta	4560
tcattgggtat aatctcacct ctttaataaa caatatgact gttgcttgta tgagcaccat	4620
taaaacgcga aatagtaacg ctttaacatc tatgattaaa aaaacctctt tcacaatttt	4680

taaaggtgca ttttaataaat agacagtatg taatcttaag aatcgaccga tgtaaataacc 4740  
 taatccattt aagaacatta atataactat caatagtcga ttttaaccata cataagacgt 4800  
 aaaatgtgca atttctaaaa atataagaat tgtgaggtat attgctaaga gtacgccaaag 4860  
 tattaaatag gtgaaataaa tccattctgt gatgtttaat ccagctaaaa agttaaattg 4920  
 aaattggttt aagtgtatga gatcggtaat catataaaat gtgtttggaa ctaataatag 4980  
 aaatatgagt ccgaaaacaa taaataaggg ccattcaaaa gctt 5024

<210> 7

<211> 9515

<212> DNA

<213> *Pseudomonas aeruginosa*

<220>

<223> Designated as P2-2

<400> 7

aagcttttct ccagaccctt caccgccttg gagatcgacg gctgggcat gtacagcttg 60  
 cgcgaggcct cggccacgct gccgcattcc acggtggtca cgaaataact gagttgcgcg 120  
 aaggtatagg acgccactgc aagacotcat cggcgcatca tcctccccgg gccgggcgtg 180  
 cgcgcctcga ttgttgtgtc cgcgcgctg caagcaagtt gcaggccgct gccgagcgtc 240  
 gcgcgctggc cgcggaacga ttgcccgcct gcacgataac ccagcacgac gcactttgcc 300  
 ggggcacgcc tggccagctt tttcttatgt cccgaggaca tttttaataa ttttcttcg 360  
 ccgcggcttg cgcgaccatc ctccccatc gaccccatgg acagcggttc gcctccccggc 420  
 ggtccggggc atgcgtgcag aaccacgacc ggcgcagacc ggcgagataa caaggagaag 480  
 gtggggtggt cgaactcagc gattggcaac ggcgcgcgcg gacacagcgc ttcacgcacc 540  
 aggccctgat cggcggcgcg cagcgtccag ccgccagcgg cgtaccttc gacgcacatg 600  
 atccggcgag caatgcctg ctggcgcggg tcggcgctg cgatgcggcc gacgtgcacg 660  
 cggcagtggc cgcgcgcgcg cgcgccttcg acgaaggccc ctgggcgcgt ctgcgcccg 720  
 tcgagcgcaa gcgcgtgctc tgcgcctggc cgagctgat ctggcccatc gcgaagagct 780  
 ggcgtgctc gactgcgtga acatgggcaa gccggtgat gacgcctgga acatcgatgt 840  
 acccggcgcc gccacgtct tcgcctggta tgcggaaagc ctgcacaagc tctacgacca 900  
 ggtcgcgcgc gccgcccagc agaccctggc caccattacc cgcgtgcgcg tgggggtgat 960  
 cggcgcggtg gtgccgtgga acttcccgtc cgacatggcc gcctggaagc tcgccccggc 1020  
 cctggccgcc ggcaactcgg tgggtgctcaa gccggccgag cagtcgcctg tctccgcct 1080  
 gcgcctggcc gagctggccc tggaggcggg ggtgcgcgaa ggcgtgctga acgtggtgcc 1140  
 gggcctcggc gagcaggccg gcaaggccct cggttgcaac ccggaggtgg acgcactgg 1200  
 gttcacccgc tcacccaggc tcggcaagta ctcatgcag tattccgcgc aatccaacct 1260

caagcaggtc	tggctggagt	gcggcggtaa	gagtccgaac	ctggtgttcg	ccgattgccg	1320
cgatcttgac	ctggcggcgg	aaaaaggcgc	cttcggcatt	ttcttcaatc	agggcgaggt	1380
ctgttcggcg	aactcgcgct	tgttggtgga	gcgttcgata	cacgacgagt	tcgtcgagcg	1440
cctgctggcc	aaggcccgcg	actggcagcc	gggcgatccg	ctggaccccg	gccagccgcg	1500
ccggcggccat	cgtcgaccgc	cggcagaccg	ccgggattct	cgccgccatc	gagcgggcgc	1560
aaggcgaggg	cgcgaccctg	ctcgcggtgg	ccgccagttg	acgatcaacg	gttcggacaa	1620
cttcatacga	ccgaccctgt	tcggcgacgt	acgcccggac	atgcagctgg	cccgcgagga	1680
aatcttcggc	ccggtgctgg	cgatcagcgc	cttcgaactc	gaggacgagg	ccatacgcct	1740
ggccaaggac	agccgctacg	gcctcggcgc	ctcgtgttgg	agcgacgacc	tgcaccgtgc	1800
gcaccgggtg	gcgcggcgct	tgaatgccgg	aacgtgtcgg	tgaataccgt	ggacgcgctg	1860
gacgtcgcgg	tgcctttcgg	cggcggcaag	cagtccggct	tcggtcgcga	cctgtcgtcg	1920
cattccttcg	acaagtacac	ccagttgaag	acgacctggg	tccagttgcg	ctgaagacgc	1980
gacggacgcg	acacgactcg	atgccgataa	cgacaacaag	aggacgatcg	aatgaacgac	2040
acgccgaacg	tgcgtgagcc	ggccctgcgc	cgcgtgctcg	ggctgggacc	gctgctggcg	2100
gtggccatcg	gcctggtggt	ttcccagggc	gtgatggtac	tgatgctgca	agggcggggg	2160
acggccggcc	tggtcttcac	cgtgccgctg	ggagtggcct	acctgctggc	gctgactacg	2220
ccttttcctt	ttccgagctg	gccctgatga	ttccccgcgc	cggtagcctg	agcagctaca	2280
cogaggtggc	catcgggcat	ttcccggcga	tcctggcgac	cttttccggc	tacgtggtgg	2340
tggcgatgtt	cgcctctcgc	gcggaaactgc	tgtgtctcga	cctgatcatc	ggcaaggtct	2400
accccggcgc	gctgccgcgc	atgctggtgc	tacggcgtgc	tcggcctggt	cacctgctc	2460
aacctgctcg	gcacgacat	cttcgcgcgc	ctgcagagcg	cgtggcgct	gctgatgatg	2520
atcgtcctgc	tgggtgctcg	cctgggtgcg	gtgagcagcg	accacgcttc	cgcgcagacc	2580
gccctggcga	gcggctggaa	cccgtggggg	gtaagcgccc	tggcgctcac	cgcgatggcc	2640
gtgtggggct	tcgtcggcgc	cgagttcgtc	tgcccgtggt	tggaggagac	gcggcgctcg	2700
gagcgcaaca	ttccgcgttc	gatgatcctc	ggcctgagca	tcattcttct	gaccatcgcc	2760
ctctactgct	tcggtgcgct	gctgtgcata	ccgcaggcgg	aactggccgg	cgacctcgctg	2820
ccacacttcc	tcttcgcca	ccgcgtgttc	ggcgagtacg	gccagctggt	cctggtgatc	2880
gccgcgatca	ccgccacctg	cagcaccctc	aactcgtcgc	tggcggcgat	cccgcggatg	2940
ctctacggga	tggcgcagaa	cggccaggcc	ttcccgcaat	tcaagcagct	cagccggcgg	3000
gcgcgcacgc	cctgggtggc	ggtgctgttc	gtcgcgcgca	tcaccggcct	gccgatcctg	3060
atcctcggcc	aggacctgga	ctcgatcaac	ctgctgctgc	tcgcgcgcgc	gctggcctgg	3120
ctgctggcct	acatcatcgc	ccacgtcgac	gtgctggccc	tgcgcgctcg	ctatccgcac	3180
atcgcctcgc	cgtttcgcac	gccgttctac	ccgctgcgcg	aactgttcgg	catcgccggg	3240
atgatctacg	cgggtggtcca	cgtctcgccg	accccgga	tgaccggacg	gatcttcgcc	3300
agcgcggggc	tgggtgctcg	cgtggtctcg	ctggtggcgg	tgggtgtggat	caagggcgctg	3360
atgcgcaagc	ccctcttcgt	acccgaaccg	ctcgagacgg	ccggtgagac	tgcccagggc	3420
aagtcgctcg	ccctcgatcc	cctgcaatcc	cttcggcctg	acgcgccaag	ggaacaagga	3480

gaacacagac	gatgaccgct	cagctcaacc	cgcagcgcga	caaccgcgac	taccagcaac	3540
tggacgccgc	gcaccacatc	caagccttcc	tcgaccagaa	ggcgctgaac	cgcgaaaggc	3600
ccgggggtga	tgggtccgcg	cgatggcctg	cagctctggg	acaacgacgg	caagcgctac	3660
ctggacggca	tgtccggcct	ctgggtgtacc	aacctcggct	acggccgcca	ggacctcgcc	3720
gccgcgcgca	gccgccagct	ggaacaactg	cgtactaca	acatgttctt	ccacaccacc	3780
caaccggcgg	tgggtggagct	ttccgagatg	ctcttcagcc	tgctgccgga	ccactacagc	3840
cacgcgatct	acaccaactc	cggtccgag	gccaacgagg	tgctgatccg	taccgtgcgg	3900
cgtactggc	agatcctcgg	caagccgcag	aagaagatca	tgatcggccg	ctggaacggc	3960
taccaaggct	cgaccctggg	cagcaaccgc	ctcggcggga	tgaagtcat	gcacgagatg	4020
ggcgcatgct	gccggacttc	gccacatcg	acgaacctta	ctggtaacgc	aacggcggcg	4080
agctgagccc	ggccgaagtt	cggtcgccgc	gcggcgctgc	aactggagga	gaagatcctc	4140
gaactgggcg	cgggagaacgt	cgccgccttc	gtcgccgagc	ccttcacagg	cgccggtggc	4200
atgatcttcc	cgccgcaaag	ctattggccg	gagatccagc	gcatctgccg	gcagtaacgac	4260
gtgctgctgt	gcgcgcgacga	agtgateggc	ggcttcggcc	gcaccggcga	atggttcgcc	4320
cacgaacact	ttcgcttcca	gccggacacc	ttgtccatcg	ccaagggcct	gacgtccggc	4380
tacatcccca	tgggcggcct	ggtactcggc	aagcgcatcg	ccgaggtgct	ggtggagcag	4440
ggcgggggtgt	tcgcccacgg	cctgacctat	tcggcccaac	cggtggcggc	ggcgggtggc	4500
atcgccaacc	tcaaggctgc	gcgacgaggg	cgtggtaacg	cgggtaaggg	aggagaccgg	4560
cccctacctg	caacgctgcc	tgcgcgaggt	cttcggcgac	catccgctgg	tcggcgaggt	4620
ccagggcgcc	ggttcgctcg	ccgcgctgca	gttcgcgcag	gacaagggtga	cccgcaagcg	4680
cttcgccaac	gagaacgata	tggcctggcg	ctgccgcacc	atcggcggct	tcgaggaggg	4740
cgtgatcatc	cgtccacccc	tcggccgcct	gatcatggcc	ccggcgctgg	tggccggggcg	4800
tgccgagatc	gacgaactga	tcgacaagac	ccgtatcgcg	gtggatcgca	ccgcgcgcga	4860
gatcggcggtg	ctctgacgcg	ccccggcggc	ccggcctcgg	ccgggtcgcc	tcgacacacgg	4920
agcgtccccc	cataacgacg	atgcggcgcc	tggcgaccgc	gcgcgggaacc	gtttcggcct	4980
ctggcgggcaa	ctgcctaagc	aacatcacaa	caatgccaat	cggtgtggg	agtgttccat	5040
gttcaagtcc	ttgcaccagt	acgcacacgt	gttttcccg	ttgtccctgt	tcgtcctggc	5100
gttcgccgcg	gcggcccagg	cgcagagcca	gagcctgacg	gtgatctcct	tcggcgggcg	5160
gaccaaggcc	gccaggaac	aggcctattt	caaacccttc	gagcgaagcg	gcggcgggca	5220
ggtggtcgcc	ggcgaatata	acggcgaaat	ggccaagggtg	aaggccatgg	tcgacgtcgg	5280
caaggtcagc	tgggacgtgg	tcgaggtgga	gagccccgaa	ctgotccgcg	gctgcgacga	5340
ggggctgttc	gaacgcctcg	accgggcgcg	tttcggcgac	cccgcgcagt	tcgtccccgg	5400
cactttcagc	gagtgcgggg	tggccacctt	cgtctggctg	atggtgatgg	cctacgactc	5460
gacgaagotg	gccagggcgc	cgcagtcctg	ggcggaatttc	tggaaagtcc	gcgagttccc	5520
ccggcaagcg	tggcctgcgc	aagggcgcca	agtacaccct	ggaagtggcg	ttgctggccg	5580
acgggggtgaa	ggcgaggagc	ctctacaagg	tactcgccac	cccggagggg	gtcagccgcg	5640
cctttcgcca	agctcgacca	gctcaagccg	aacatccagt	ggtgggaggc	cggcgcccag	5700

ccgccgcaat	ggctggcggc	cggcgacgtg	gtgatgagcg	cggcctacaa	cgggcgcatc	5760
gcgcgtgcgc	agaaggaggg	ggtgaaactg	gccatcgtct	ggcccggcag	tctctacgat	5820
ccggagtact	gggcggtggt	gaagggcacc	ccgaacaagg	cgctggcgga	gaaattcatt	5880
gccttcgcca	gccagccgca	gacgcagaag	gtgttctccg	agcagatccc	ctacgggccc	5940
gtacacaagg	gcacccctggc	gttgctgccc	aagacgggtgc	aggaggcgct	gccgacccgc	6000
gccggccaac	ctcgaaggcg	cgcgggcggt	ggatgccgag	ttctgggtgg	accacggcga	6060
ggagctggaa	cagcgtttca	atgcctgggc	gcgcgctgag	cgctgcgcgt	cggcaaaaaa	6120
aatgacgggc	cccaagtcgt	ccgggcccgt	cgggtcaaag	cgctgacggg	gtgatcagcg	6180
cagctcttcc	aacaacccct	gcagataccg	acagccctcg	gtatccagcg	cctgcacccg	6240
aaggcgccgc	gccccacact	ccaggccgga	gaggcccgag	ccggccttga	tgggtggtcg	6300
caggccccgg	cggaggatga	agtcgagcag	cggcaactgc	cggtagaaca	gcgcgcgggc	6360
cttctccagg	tcgcgcgtga	gcaccgcctg	gtagagctgg	ccgttgagcg	tcgggatcag	6420
gttcggcgcg	gcgctgcacc	agcctttcgc	gccggccaac	aaggcctcca	gcgccagcgc	6480
gttgacgccc	ttgtagaagg	gcacccggcc	ttcgccgagc	aggcgagct	tgtgcatgcg	6540
ctggatgtcg	ccggtgctct	ccttgaccat	ggtcacgttg	tccacttcgc	ggacgatgcg	6600
caggatcagt	tccaccgaca	tgtcgatgcc	gctggtgccc	gggttggtgt	agagcatcac	6660
cggcacgccc	atggcttcgc	caaccgcgcg	gtagtgcctg	aacacttcgc	cctcggttgc	6720
cttcacgtag	gagatcggca	ggaccatcac	cgcctcggcg	ccgagggatt	cggcgaaactg	6780
cgcgcggcgc	acggtcttgg	cgggtggtcag	gtcgggagac	ctgacgatgg	tcggcacgcg	6840
atgggcgacg	gtcttcaggg	tgaagtogac	caactcgctc	cattccgggt	cgtcaggta	6900
ggcgcccttc	ccggtgctgc	cgcgcggggc	gatggcgtgc	acgcgcgcgt	cgatcaggcg	6960
ctcgatggag	cggccgaggg	ccggcaggtc	gagacgcgcg	tcggcgccga	aggggggtga	7020
tgggtgtagc	gatgatgccg	tggatggatg	cggacattgg	atgtaccctg	gacattgagt	7080
gggaaatgcc	aggacggacc	tgggtgggaa	ggtcgcttcag	ctcaggcagt	cgtcgttgcg	7140
cggcaggcag	cgcgcggcgt	agtagttgaa	tgcggcgccg	tggcgcttcg	gggtggagat	7200
ccagtcgtgg	gcctcgcgcg	ccagggcggg	cgggatoggc	ttgatctctc	cggcgggccat	7260
cgcacgcaac	tgcattcttc	ccgcgcgcct	gagcagcacc	gcgatcacgc	aggcctctct	7320
gatgctcgca	ccggtggcca	gcaggccgtg	gtgggagagc	aggatggcgc	gcttgctgcc	7380
gaggcgggcg	gagatgatct	cgccttcctc	gttgccctacc	ggcacgcccg	gccagtcctt	7440
gagggaaggcg	cagtcgtcgt	atagcgggca	aaggctccatg	tgcgagacct	gcagcgggtac	7500
ttccagggtc	gacagcgcg	cgatgtgcag	cgggtgggtg	tggatgatgc	agttgacgtc	7560
cgggcggggcg	cgatagaccc	agctgtggaa	gcgattggcc	ggattcgcca	tgcgctgccc	7620
gtggaggacg	ttgaggtctt	cgtcgaccag	cagcagggtg	ccggcgctga	tctcgtogaa	7680
gcccaggccc	agttgctggg	tgtagtaggt	ccccgcctcc	gggcgcgcgc	aggtgatctg	7740
ccggcgagcg	ccggagtcgt	ggccggccctc	gaagagaatc	cggcagggtca	gggccagctt	7800
ttgccggtca	gtccacgtat	tatcgccgag	gctgcttttc	atctgcttca	gcgcgtgctg	7860
gatcagttga	tccttgggta	attccagtgt	cgtaaccatg	cgaggttccct	ttgacggagc	7920



```

gagtcggggg aaacggccagg cagttgcgcg ccacgcaacg acccggtgt aaatgacacg 7980
gatcaagtta tatgacacaa agtgtcattt agcaagagag aagtttcacg gccatcggga 8040
gaaggctgtc ctcaatgtcc atgcgcttga aattgctgag aaaaaaactc ggggtcacgc 8100
tgagagaccct ggccgacaag accggcctga ccaagagcta cctgtccaag gtcgagcgcg 8160
ggctgaacac gcggtccatt gccgcgcgcg tgaagctggc gaaggcggtg aacgtgcagg 8220
tgaggaggct gttctccgag gaaagcgacg gtgtcgacgg ctacagcatc gttcgtcgcg 8280
accagcgcaa gtcgctgtcc agcggcgacg acggcccggc ctacgcctcc ctcgtcgcag 8340
cagatcggcg ccgcgcgcgt gttgcggttc atcgtccacc cccgcgcga tttcagtcac 8400
tcgacgttca aggagcacct cggcgaagag ttcatcttcg tccatgaggg ccaggctcag 8460
gtcgacttca tgaaccagcg gatcatcctc gagcgcggcg acgccttgca tttcaacgca 8520
cagaagccgc accgcacccg ctccctgggg gagaccaggg cggaattgct ggtggtgatc 8580
cacagcgacg aatgaggcga cggcttcggt cgatcggatg ctgctaacg ttctgttcga 8640
ttatcgaaat gttaatgat tatcggaattg tgagccctcg gaccccgcg taaggttctc 8700
gtcacgtgcc gtccaggcag cgcacaacaa gacgagaccg gaccgatggc tgaaatcctc 8760
tccttgcgcg aacgggtgca cgcttcgtcc acgatggcga cagcgtcgcc ctcgagggt 8820
tcaactacct gatcccgacg nccgcgggc acgagctgat ccgccagggc aggaagacc 8880
tgacgctgat ccgatgact ccgacctgg tctacgacct gctgatcggg gcaggctgcg 8940
cgaagaagct ggtgttctcc tggggcgga accccgggtg cggttcgctg caccgcctgc 9000
gcgacgcggg ggagaagggc tcggccgcaa ccgctggaga tcgaggaaca cagccacgcc 9060
gacctcgcca acgcctatct tgcggcgcc tcgggctgc ccttcgcggt ntgcgcgct 9120
acgcgggctc cgacctgccg aagggtcaacc cgctgatccg cagcgtcacc tgcccgttca 9180
ccggcgaagt gctggcgcg gtgccctcgg tgctccgga cgtcagcgtg atccacgcgc 9240
agaaggccga ccgcaagggc aacgtgctgc tctggggcat cctcggcgtg cagaaggaag 9300
cggccctggc ggogaagcgc tgcatcgta ccgtcgagga gatcgtcgac gaactggacg 9360
ccccgatgaa cgctgcgtc ctgcgagct ggggcgctca gcgcggtgt cctggtgcc 9420
ggcggcgcg atcgtccta tgcccacggc tactacgagc gcgacaaccg ctctaccag 9480
gactgggacc cgatcgcccg cgaccgcgaa agctt 9515

```

<210> 8

<211> 2291

<212> DNA

<213> *Enterococcus faecalis*

<220>

<223> Designated as EF-1

<400> 8

aagctttaga	taatgataaa	cgcggtgtatg	tgaatgtcca	gccgattcaa	tcgcctactg	60
gagaaacagt	gattgggtgtc	ctttatgtga	aaagtaattt	agaaaataaa	taccaagaaa	120
ttactaacac	agcaagtatc	tttttcaactg	cttctattat	tgccgcagca	atctcgatta	180
ttgtgaccct	actgattgca	cgatcaatca	cgaagccgat	tggtgaaatg	cgcgagcaag	240
ccattcgaat	cgctcgtggt	gattacgctg	gaaaagtaga	agtcocatgga	aaagatgaat	300
taggcccaatt	agcagaaaaca	tttaatacaat	tatcagaacg	gattgaagaa	gcacaagaaa	360
caatggaagc	agaagaatcg	tttagatagt	gtcttaacgc	atatgacaga	tggtgtcatt	420
gcgacggatc	gccgcggaaa	ggtgattacg	attaatgaga	tggtccctttc	attattaaat	480
gtaaaaaatg	aaaatgtgat	tgggacctcg	ttattagagt	tgttagatat	tgaagaagat	540
tacacattgc	ggaagctggt	agaagagcca	gatgaactgc	tgattgatcg	ctcaacgtct	600
gacgtgaag	aagaccaaatt	gattatccgg	gtagacttta	cgatgattcg	tcgggaatca	660
ggatttatta	ctggcttagt	ttgcgtactt	catgacgtca	cagaacagga	aaaaaacgaa	720
cgggaaagac	gggaatttgt	ttccaatggt	tctcatgagt	tgcgacgcct	ttgacaagta	780
tgcgtagtta	tatagaggct	ttgagtgaag	gagcttgga	aaacctgag	attgcgccga	840
atttcttaaa	agtcacgtta	gaagaaaccg	accggatgat	tcgtatgatt	aatgatttgt	900
taaatttatc	tcggatggac	tctgggaata	cacatcttca	attagagtat	gtgaatttta	960
acgaattgat	taattttgtc	ttggatcgct	ttgatatgat	gattgaaaat	gagcaaaaaa	1020
attacaaaat	tcgccgtgaa	tttactaaac	gcgatttatg	ggtagagtta	gatacagaca	1080
aagtaattca	ggtttttgac	aacattttga	acaatgcgat	taagtattcg	ccagatggcg	1140
gcgtcattac	ctgccgacta	gttgaaacac	ataataatgt	cgtcttttagt	atctcggacc	1200
aaggtttggg	catccctaaa	aaagatctcg	ggaaagtott	cgagcgtttt	tatcgtgtgg	1260
ataaagcacg	tcgcgcgagca	caagggtggga	ctggtttagg	tttagcaatt	tctaaagaag	1320
taattcgggc	ccataacggg	agtatttggg	tggaaagtac	agaaggtgaa	ggatcaactt	1380
tctatatattc	actaccatat	gaaccttatg	aagaggattg	gtgggaatga	tgaaaaaatc	1440
agaatggatt	acaagaattg	gcttgatttt	gatgggtcatt	ttaagtatat	atttttcaagt	1500
caatatctgg	ctgaattctg	ccaaaaaat	accagaaatg	aagtcgggaa	gccaaagtcac	1560
aacagctgtc	aatgaaaaag	ccattggcga	tgtctatttta	cctttgcaat	tgattcgaat	1620
agccgatgga	aaagcgatgc	aaagtaatcg	tgaaacatta	attagtaatg	ttcaaaatga	1680
tattaaaatg	gctacgtttg	gtaaattgac	acaagttgtg	acaaaaaatg	cagagcaact	1740
taagcgctac	aaccaaattg	aacaaggcat	tgaacttctt	tatcaaggtc	cctttttaat	1800
ctcggactat	gcttcgattt	ataatctatc	cattaatttt	actaacttta	atgagttgac	1860
ggaccagtat	tttacgaaaa	ttcaattgga	ttttaacgaa	aataagatac	gttttttaga	1920
ttatgatcaa	tccaacgtct	atgaagcgcc	catgactggt	aataaggcgc	gcttaatggg	1980
aattatcaat	aaagagggat	tgcaatatca	agacgtttcc	gaaaatacgc	taaccaaaca	2040
aggacaatgt	tatttaacca	atgatatgaa	gttgaaaaag	tacagttata	tottanttcg	2100
caaccagtta	ctcgttttag	gaatgctttt	ttcaatgaaa	cggaagatat	ccaaaccaat	2160
gaagacagtc	aagacttaac	ctatacgagt	aaagaagaac	gattgtttgc	agaagaaaaa	2220

ctggggaaaa tcgattttta agggaaccttg ccagaagaga ataaacggga ctcaatctat 2280  
aatcaaagct t 2291

<210> 9

<211> 2441

<212> DNA

<213> *Enterococcus faecalis*

<220>

<223> Designated as EF-27

<400> 9

aagcttctgc gctaggaacc agccctttta ttacatctcc ccatactgga tttgacaatg 60  
ccacttgata agcaaaaatc acaaaaataa caacaattaa agcaacaaca atagcttcaa 120  
tttttctaaa accaattttt gtcaataaca acaaaagtaa aacatcaa ataccgtaatga 180  
agacagccag acctaaagga atatgaaata ataaatataa ggcaattgcg ccccccataa 240  
cttcagcgat atctgtagcc ataattgcta actctgttaa aatccataat acaataccta 300  
acgtcttact agttctagca cgaatcgctt gtgctaaatc catctgtgaa caatgcctaa 360  
tttagcagcc atatattgga gcaacattgc aatcaaactg gaaattaaaa taatcgacat 420  
caataaatat tgaaaatttt gtccccagc aattgaagta gaccagtttc ctggatccat 480  
atacccact gctaccaatg ctcttgacc tgagtaagca aataacgttt tccaaaaact 540  
catattttta ggcacgtcga tgggtgccatt aatttcttca agcgaaggac catttgcata 600  
ttcaatcaaa tgatgtcttt gctttggttc atgttcttct gaatttttca attcaattcc 660  
ttctttcggt ttgcaataat tttaaaaggc ccttcccgtt agaagggtta cctctagtat 720  
attttaggta cacctaaaat atactgctaa aaataacaaa atgcaagact tgaaagaaaa 780  
ttttgacagt gtaaaaatag attgtcgtaa atgtgcgac ttaaagtttg aagaaatcag 840  
ggtagctggg agttgattat cttaagaagt agaaaataag ggacctaagt catttcggct 900  
taggtccctt attttatttt tattcggtta ttctattaag aatggatgct acaatttctg 960  
tcgtgtcagc tgaatgattt ctaaaatctc gtaaaactta tctgacgaaa accttcaagt 1020  
acttcgggca acttattttt cccccattca aaagttccat catttctttt caataatctt 1080  
tgtaaaattt cttctttctc gaccgctaac aaaaaatgat aaacgtcaat gcctgctcgt 1140  
ctcagatata caatcagctc ttcttcata tcatTTTTat aaagggtcat tgtaacaata 1200  
atcgcccgtc cagactcttt ggacattcgt ttttaataat gagcattcca gcaacgccat 1260  
tcttgatact cctgaaaatc attttctttc atttcttcgg gaactagctc catcaatgca 1320  
ctaccaataa tttctggatc ataaatgatt gcgttgggaa gtttttggtg taactcatgt 1380  
gcaatggctg tttttccgga tccaaacgca ccgtttaacc aaataattat cataatttcc 1440  
ttttcttctg aacaaatttc tttgttgttt aatttaggtg ctagattaact ttttaattttt 1500

```

ttagccattc acttatagtt actacttaca tctttaacag taaacgagac aaactaaaaa 1560
tacaacatcc tacgctatta acctcgggtt atataacata ctcatctgat aatttctccc 1620
taaaaaaaca gaatgtgggc aatcttttta agaataattg aatagaataa caacaaacag 1680
taattcaggt ataaccagct agaaattggt ttatttttag tcacgagtat gataagcatg 1740
taaatcaaat agaatcatat taggtgaggt tactctgaag aacacagggt atcgctcgga 1800
aatgtcgaga gacagtaacg agtaaagcag ggattgtcga attaaggctt tcctaagata 1860
actagaattt ttttcttacg tctcagaaaag ccaaagctca attattgtga ttaccctata 1920
atcttcttct tttattcggc gacctcttta atatgattaa ttggagggtt ttaaattgaa 1980
agctgtcact gcatcatcta agaaaaatac cctacttgct aaaagtatcg ggaatcttac 2040
cttgtctcatc attttaggca ttttcatttt tatcatcgtc ttctcttggc taaaaatgaa 2100
tcgccctctc cacacccttc cctcagaaga attcctcgca acaccaagta aaacagatga 2160
tttcttatct ccatcaaatc ttttttactt ttcaattcga accatgtttc gaatgattgt 2220
ggggatggct tggtccttcc tgttttcctt tgtttttggt atttttagccg taaaatataa 2280
aacggcacga agagtcattt taccattagt taatttcctt gaatctgttc cattgctagg 2340
ttttttgacc ttacaaactg cttggttact tggtttattt ccaggaaatg tgatgggcgc 2400
agaagcgggt gctatttttg ccatcttcac aggtcaagct t 2441

```

<210> 10

<211> 3480

<212> DNA

<213> *Enterococcus faecalis*

<220>

<223> Designated as EF-7

<400> 10

```

aagcttctag cgtttcggat tggcgccctat gatgcaccag gagagcgacg aatcaatacc 60
aaaaatatgc ctacagcagg aggacttgca atctacattg cttttgctag ttcatgttta 120
ttgatttttc gttcgattat ccacaaagat tatatttggc cgattatttt ggctggtgga 180
atgggttgtt tgacaggcct cattgatgat attaaagaga ttactccaat gaaaaaaca 240
atcggatatt tgtagcagc attagttatt ttattttggt gctggaattc ggatagattt 300
tgtgacgttg ccagttgttg gaatgattga tttgcgctgg tttagtttac cactaacttt 360
attgtggatt ttagcgatta cgaatgcagt aaatttaatt gatggtttgg atggtttagc 420
atcaggcgta tccattattg gattaaccac gattggtatt acagggtatt ttttcctaca 480
tgctaaaacg gtctatatcc caattgttat tttattttta gttgcgagca ttgcgggatt 540
tttcccatatc aatttttatc cggctaaaat atttctagga gataccgggg cgttattcct 600
cgggtttatg attgcagtaa tgtcgttaca gggcttgaaa aatgctacgt ttattacggt 660

```

aattacgcca atggtgattt taggtgtgca attacggata cggtttatgc aattattoga	720
cggctattga acaagaagcc catttcctca gcagataaaa tgcattttaca tcaccgcttg	780
ttatcttttag gttttaccca taaaggggog gtcattgacta tttatgcatt agcggttagtt	840
ttttcctttg tctctttatt gttcagctat tcaagtacag tagcatcaat tttattaatt	900
gtcttttgtt taattggctt agaactatctc attgaactaa tcgggtctagt tggcgaaggg	960
catcaaccgt tgatgtattt gttacggatt ttagggaatc gtgaatatcg tcaggagcaa	1020
atgaaaaagc gacttggcaa gcattctaa agaaagtaaa gaaatcttta ggttgctttg	1080
cgaagagctaa acctatgata taattccatt aaacttaaaa aagtatatgt gtgaaacata	1140
tgtttttttt ttaagacgat gtttcagtag taaggagaaa tgagcatgca agaaatggta	1200
acaatctoga ttgtcactta taatagtcgt tacattttta atgtactaga ccaattaaaa	1260
gccgaactag gtactgatag tatctatgat attcatatct atgacaatca ttctgaaaca	1320
gcgtatcttg aaaaattaac aacatatgaa ccatttatta ctatccatcg cgctgaagaa	1380
aatcaagggg ttgggtcatgg tcataatcaa gtgttattca atgcttcgac aaagtatgca	1440
attattttta tcccgatgtg ttggttacta aagacgtgct tgatogttat tagacgtatc	1500
aaatagataa gaacattgca gtccgttagcc cttaaagttgt taaatgaaga tggcacgacg	1560
caatatttag ttctgcaaaa attagatgtc ttcgattata tgttaogttt tattcccttt	1620
caatttgtaa agaaaatttt tgataaacgt ttgagtattt atgaatgtcg cgatttgctg	1680
gatacagaaa caacggatat taaaatgggc tcaggctgtt ttatgttgat tgatcgtgaa	1740
aaattcgttg aaattgggtg gttcgatgaa cgtttcttca tgtactttga agacaacgat	1800
ttatgtttac gctttggcaa agcaggctat cggattctct atacgccttt tgaaacgggt	1860
gttcacatgt atgaaaaggg cgcacataaa agtcgaaaat tgtttaaaat ctttatgcaa	1920
tcaatgggga aattttttta caaatggggc tggaggttct tttaatgagt caaagattag	1980
cggtagtcat cgtcttatat caaatgaaaa tggctgatac gccgaattat ttgttattaa	2040
aagaagtggg agaccacccc caattgcact tatttattta tgacaacagt ccacttcctc	2100
aagaagatgc attattttta caaccaaatg ttacttatcg acataatcct gataatccag	2160
gactagcgac cgtttataat gaagcgattg ctttttagtca agcgaatcaa tgtgaattat	2220
tggtgctcct tgaccaagac acagaagtgc cagcctotta ttttgatacg ttgatcatca	2280
tgccattaga tccgactgtg gcagtctatg ttccaattgt agaagcaa at ggacaacaaa	2340
tttcgccagt atatagtgat caatacgttg ggcttaaagg agcaaagcca acagcaggga	2400
tagccaacca accgttgatg gctatcaatt ctggtacagt tattacggca gaaacgctac	2460
gctgggttga aggattttcg gaagaatttc ctttggaacta ttttagaccat tggttctttt	2520
atcaattaaa tcaagccaat aaaaagattg aagtcttacc aatccacctt aaacaagaat	2580
tgtctgtttt agattatcgt acaatgagtc ctcaacgtta tcgctctatt attgaagcag	2640
aaacgttatt ttatcgtcga tatgatcaag aaaagtttcc ccatcatcga cgccatttat	2700
ttttacgcag tagtaagcaa tttttaactg tcaaaaatcg ccaaatttgg cggcaaacat	2760
tggcagaatt tctcaagtta atgaaaggat aatctatgat ctcagtttgt attgcgacat	2820
ataatggaga aaaatatctc gcggaacaat tagatagtat tctttttacaa gtcagtgaag	2880

```

aagatgaact aattatttca gatgatgggt ctactgatca tacgttggaa attttgagga 2940
cgtatgcagc gaattatccc caaattcaat tgttacaagg tcccagggca aggagtgatt 3000
gctaattttg ctttttgcct tacgcatacg aaaggcgaag taatattttt agcagatcaa 3060
gatgatgttt ggttgccaaa taaagtaacg acggtgacag aatattttga agcgcaccct 3120
gacatccaag tggttattag tgacttgaaa attgttgatg cggatttaca agttaccaat 3180
ccctcttatt taagtttcca aaagtcaaac cagggttttg gcgaaatgcg ataaaaagtg 3240
gctatatattg ggcaggtatg gcctttcgtc aagaaatgaa aaacgtcatt ttaccattc 3300
cgccagaagt tcctatgcat gatatgtgga ttggcttatt agctgcacgg aagaagcaaa 3360
cgggtctcat taaagaacca ttagtgcttt accgaagaca tggagcgaat gtcagcccca 3420
ttattaccaa aacaagtttc caacaaaaat taaattggcg tgtgaattta ttaaaagctt 3480

```

<210> 11

<211> 3615

<212> DNA

<213> Escherichia coli

<220>

<223> Designated as EC-24

<400> 11

```

aagctttttct tgcgtgttct tgtgaggctt oottogccat tatcatcacg atccacataa 60
ataaagccgt agcgccttaga catttgtgaa tgagatgcac tgactaaatc aattggcccc 120
caactgggtgt accccataat atccacacca toggcaatcg cttcatttac ctgtaccagg 180
tgatcgttta aataggcaat tcgataatcg tcctgtatcg aaccatccgc tccaacgctg 240
tcttttgcgc ctaatccgtt ctgcacaata aataacgggt tttgataacg atcccaaagc 300
gtatttaaca gaaccogtaa tccaaccgga tcaatttgcc acccccactc tgaacttttc 360
agatgcggat tggggatcat attcagtatg ttgccctgcg catttttatt aatgctttcg 420
tcgtgggaac acaaccagtc atgtataact aaagagatga atcgacggta tgttttaaat 480
ctctgcgtca ctttcagtca totcaatggt gatattgtgg tcogcggaaga aacgctgcat 540
atagccggga tactggccac gcgcctgaac atcaccaaag aacatccagc gcgggttctc 600
ttccatggcc tgcaacatat cctgtggctg gcagggtgagg gggtaaacca gccacccag 660
aagcatattg ccgattttcg cttcggggag caggctatga caggctttta ctgcccgccg 720
actggcaacc agttgatggt ggatagcctg ataaacttcc gcctcgccac tctctttctg 780
cagccccacg ccggtgaatg gcgcgtgtaa cgacatgttg atttcattaa acgtcagcca 840
taacgccact ttatgttggt agcgagtaaa gaccgtgcgg gcgtaatgtt cgaagtgate 900
gatgaccgt cgattagcca accgcgtag tttttacca gcccatatgg catttcgtaa 960
tgggataacg ttaccagcgg cttgatcccc gcctgcgcca tttcatcaaa cagccgatcg 1020

```

taaaacgcta	accccgcttc	attcggttcg	acttcgtcgc	cctgagggaa	aattcgcgcc	1080
caggcaatgg	aaatacgcag	acaggtgaag	cccatctcgg	caaataacgc	gatatcttcc	1140
gggtaacggt	gataaaaatc	gatggcgaca	tctttgatat	tctctttccc	caggatgcgc	1200
ggttccattt	ttcccattac	gcatgaggct	gtaaatctga	ggtcgagatc	cctttgccat	1260
cttctgccca	ggcaccttcc	acctgattgg	cagctgttgc	ggcaccccaa	agaaatgttt	1320
ctggaaatgc	tttcataatt	aactcctttt	atcgtttagcg	aatgatggat	aacagcgggt	1380
cacctgcgct	tatctgcgcc	gtgccgtggg	gtaatacgtc	cgtaaaatca	tcgctattac	1440
tgattaatac	cggcgctcgc	agatcaaata	cggcctcgcg	aatagcaggg	atatcaaaag	1500
aaatcagccg	atcgctgta	ttgaccttgt	caccacggtt	gacgtgagcg	gaaaagaatt	1560
tgccgtccag	ttttacggtg	tcgataccga	catgaatcag	gatctccaca	ccatcatctg	1620
actcaatgcc	aatggcgtgt	aatgtggcga	acaacgaagc	aattcgaccc	gcaaccggag	1680
aacgcacttc	accaaccgag	ggcagaatgg	caataccttt	acccaacagg	ccactggcaa	1740
acgtgggtatc	agcgacgtga	atgagcgaca	caatctctcc	cgtcatcggg	gaacagatac	1800
cgccttgcctc	aggtggtgta	ataacctctg	gtgttttctc	ttcggggcac	cctgcgctgg	1860
ctgacgttta	gcggtgatga	aatgaagcat	caccgtaccg	acaaatgcgc	aaccgatggc	1920
aatgacaccg	ccaataacgc	tggcccagac	ggtgaaatca	attcccgttg	acgggatggt	1980
ttgcatgaag	gtgaaaatac	ttggcaaacc	aaaggagtag	actttcgttt	gcgcgtagcc	2040
aataatgggtg	gccccaaaag	ccccactgat	acaggcgata	acaaaggggt	acttacgcgg	2100
caggttgacg	ccatataaccg	ctggttcggt	gataccaaac	agactcgtca	acgccgctga	2160
tcccgccacc	acttttttct	gcgcctcgcg	ttcgcagagg	aagacgccga	gcgcgcgcgc	2220
gacctgcgcc	ataatggcgg	gcattaacag	cgggatcatg	gtgtcgtagc	ccagcacggt	2280
gaagtatttg	atacacaccg	gcaccaggcc	ccagtgcagt	ccgaacatga	cgaagatttg	2340
ccagaagccg	cccattaccg	cgcccgcaaa	tgcaggaaac	gcctgataaa	gccagagata	2400
accggcggca	atcagttcgc	ttatccaggt	tgatagcggc	cccaccagca	gaaagggtgac	2460
gggtgtgata	accatcagac	atagcaatgg	tgtgaagaaa	tttttgattg	ccgacggtaa	2520
ccacgcatta	agtcggcggt	ccagaatgct	gcacaaccag	gcagaaaaaa	taatgggaat	2580
aaccgatgac	gagtaattca	acaatgtgac	cgggaataccc	aggaaatcca	gccccagcgc	2640
atccgccttt	gcgcgttctc	gaaaagcagt	acagaattaa	tggatgcact	aacgcctccac	2700
caatcaccat	ggcagtaaat	ggattatcgc	cgaagcggtt	ccccgcggtg	tatcccagga	2760
ttatcgggaa	gaacccaaaac	aaggcatcac	tggcgtgtaa	taaaattaaa	taagtaccac	2820
tttgttcggg	cgtccactga	aaagtgagcg	ccagagccag	catacctttc	aagatccccg	2880
gttgcccgcc	atcaaaccga	tacagaggcg	taaaaatacc	tgaaataaca	taaacaaaagc	2940
ggtttagaca	gattaccttt	atcatacatt	ttccggtgcc	tgttgcgctt	tttcgtcaag	3000
gcctgccaca	ctgttaaccg	ccaggaagac	atcggccaca	tggttaccta	tgaccacctg	3060
aaactggcca	ccgctttcca	ccaccataat	aataccgggg	gtctttttca	gtacctctgc	3120
ttgcgccttg	ctttcatcct	ttaattttaaa	aacgtaaaac	gcgttgcgca	atgcatcaga	3180
ctcacaatgt	tatctgcgcc	cccgaactcct	gcgactatit	ttctggctaa	ctccgtcata	3240

```

acttgccctc tacgctttgc ggcaaaactc caaaaaaaaaa cctgaaaaaa acggcctgac 3300
gtgaatcaag caattttttt caggttttgc cgccttagtg cggtaacaat cctttactca 3360
gtaataatat ttcagtgttc tttgcgcacg cgcctctatat ttatggctaa aaacataatc 3420
tctgcggtg aaattttacg ttgatactgc aaaccaataa aaatggcgat ccgttccgca 3480
cattgccatg cttgcgggta attttgtttt actgcttggt gtaatgattc atcaactatc 3540
ttaattgaag catgttcaag aatagccag gataaaaact tcagatgtgt aaccagtcgc 3600
tgataactca agctt 3615

```

<210> 12

<211> 4954

<212> DNA

<213> *Escherichia coli*

<220>

<223> Designated as EC-34

<400> 12

```

aagcttaacc gctctcatct gttgacgcga cggcatagct atattctgcc ggtcctggga 60
cgtagcgaga ttgacatgca aaaaaacggt ggcgaggcgg taaccgttga ggattcaatg 120
togatgattc atgcctcgcg tggcgtgtta aaaccgcgcg gtgtaatgct gaaatcagag 180
tgtgcagtggt tcgcgggaat cgcgcaggca gcactacccc agagcgtggt agcctgggag 240
tatctggtgg aagattatga tcgcattcgc aatgacattg aagctgtgct gccagagttc 300
gccgactata accagcgcgt ccgtcatccc ggtgggtttc acctgataaa tgcagctgct 360
gaaaggcgct ggatgacgcc gtcaggtaag gctaatttca ttaccagcaa agggctgtta 420
gaagatccct cttcagcgtt taacagtaag ctggtcatgg cgacagtacg cagccacgat 480
cagtacaaca cgacgattta tggatatgat gatcgctatc gaggggtatt cgggtcaacga 540
gatgtggtct ttatgagtgc taaacaagct aaaatttgcc gtgtaaaaaa cggcgaaaga 600
gttaatctta ttgcgcttac gccagacggt aagcgcagtc acgcgcgatg gatagattaa 660
aagtggtoat ttaccctatg gctgaccgct cactggtgac ctattttcca gaatcgaatc 720
acatgctaac acttgataac cagcatccat taagtggcat tccggtctat aaaagtattc 780
cgcttgaatt agaaccatca aattaatgtc tcttctcatt tcttctgctg tcatccgcac 840
agcagaagaa ttcctcattg actattatct cgcaatttgc tcacatggat taaattaaac 900
tacatactat aagatataaa cttctgccta cagctgtaag aaactccgct cagtactgaa 960
gcaccagtcc tatttcctct tttctccagc ctgttatatt aagcactatg attaacgatt 1020
tttaacgtta tccgctaaat aaacatattt gaaatgcatg cgaccacagt gaaaaacaaa 1080
atcacgcaaa gagacaacta taaagaaatc atgtctgcaa ttgtgggtgt cttattactg 1140
acacttaagt gatagccatt ttttcggcaa ttgatcagct gagtatttca gaaatgggtc 1200

```



gcattgcaag agatccttaca catttcatta tcaatagttt gcaaggctgt aaacaaacag 1260  
caaattataa atatgaaatg ttaaaaaagt atcgataaaa actttattgt ttttaaggaga 1320  
taaaatgtcg ctcgtttggt ctggttatatt tattcatcat gccctcaacg ctaacatttt 1380  
agataaagat tacgccttct ctgacggcga gatcctgatg gtagataacg ctgttcgtac 1440  
gcattttgaa ccttatgagc ggcattttta agagatcgga ttactgaaa ataccattaa 1500  
aaaatatcta caatgcacta acatccagac agtgacggtg cctgttcctg cgaagttttt 1560  
acgtgcttca aatgtaccga ctggattgct taatgaaatg attgcttata tcaactcgga 1620  
agaacgcaat catcataatt ttccagaact ttgtctttt tottgctgt ctatttttgc 1680  
cgcatgcaaa ggtttcatta cactattaac taacggtgtg ctatccgttt ctgggaaagt 1740  
gagaaatatt gtcaacatga agccggcgca cccatggaag ctgaaagata ttgtgactg 1800  
cctgtacatc agtgaaagcc tgttgaagaa aaacttaagc aagagcaaac gacattctca 1860  
cagattcttt tagatgcaag aatgcagcac gcaaaaaatt tgatacgcgt agaaggttca 1920  
gtcaataaaa ttgccgaaca atgtgggttat gccagtacat cttattttat ttatgcgttc 1980  
cgcaaacatt tcggcaacag tccgaagaga gtttctaagg agtaccgttg tcaaagtcac 2040  
acgggtatga atacgggcaa cacgatgaat gctttagcta ttgattatt tgctaacgag 2100  
tagtcaacca cacacgctgc gtaagaatta aatggggcag ccattccctg ccccgcggtg 2160  
tttttaggcg atatatttat tgaaataaat aagtgaacac catcacatat ttatgcactt 2220  
gcataacctg ttgcatgatt atttatgatc tcaattctgc attttgtcag taaaatgcaa 2280  
taatttatta aatatcaata aattagttgt ttatcggcga gaaattactt aatagaacag 2340  
aaagtaatgt caacgcttta tggactgttt ttccctttt tttagctaaa totgctatct 2400  
ctttatgtga ctaacttcac ttacatccac ttatttctct tcgtaaaatt actttggaat 2460  
taagtacaat aagaagagga acatttatga agtctgcatt aaagaaaagt gtcgtaaagta 2520  
cctcgatata ttgtactg gcactggta tggctgcatt tgctgctcat gcggcagatg 2580  
atgtaaagct gaaagcaacc aaaacaaacg ttgctttctc agactttacg ccgacagaat 2640  
acagtaccaa aggaaagcca aatattatcg tactgaccat ggatgatctt gggtatggac 2700  
aacttccttt tgataaggga tcttttgacc caaaaacaat ggaaaatcgt gaagttgtcg 2760  
atacctacaa aatagggata gataaagcca ttgaagctgc acaaaaatca acgcccagcg 2820  
tcctttcatt aatggatgaa ggcgtacgtt ttactaacgg ctatgtggca cacgggtgtt 2880  
ccggccctc ccgcgccgca ataatgaccg gtcgagctcc cgcccgcttt ggtgtctatt 2940  
ccaataccga tgctcaggat ggtattccgc taacagaaac ttcttgcct gaattattcc 3000  
agaatcatgg ttattacact gcagcagtag gtaaatggca ctgtcaaaa atcagtaatg 3060  
tgccgggtacc ggaagataaa caaacgcgtg actatcatga caccttcacc acattttctg 3120  
cggaagaatg gcaacctcaa aaccgtggct ttgattactt tatgggattc cacgctgcag 3180  
gaacggcata ttacaactcc ccttcactgt tcaaaaatcg tgaacgtgtc cccgcaaaag 3240  
gttatatcag cgatcagtta accgatgagg caattggcgt tgttgatcgt gccaaaacac 3300  
ttgaccagcc ttttatgctt tacctggctt ataatgctcc gcacctgcca aatgataatc 3360  
ctgcaccgga tcaatatcag aagcaattta ataccggtag tcaaacagca gataactact 3420

```

acgcttccgt ttattctggt gatcagggtg taaaacgcat tctogaacaa ctgaagaaaa 3480
acggacagta tgacaatata attattctct ttacctccga taatgggtgg gttatcgatg 3540
gtcctctgcc gctgaacggg ggcacaaaag gctataagag tcagacctat cctgggggta 3600
ctcacacccc aatgtttatg tgggtggagaa ggaaaacttc aaccgggtaa ttatgacaag 3660
ctgatttccg caatggattt ctaccgcaca gctcttgatg cagccgatat cagcattcca 3720
aaagacctta agctggatgg cgtttccctg ctgccctggg tgcaagataa gaaacaaggc 3780
gagccacata aaaatctgac ctggataaac tcttattctc actgggttga cgaggaaaat 3840
attccattct gggataatta ccacaaattt gttcgccata cagtcagacg attaccgcga 3900
taaccccaac actgaggact taagccaatt ctcttatacg gtgagaaata acgattatct 3960
gcttgtctat acagtagaaa acaatcagtt aggtctctac aaactgacgg atctacagca 4020
aaaagataac cttgccgcgc ccaatccgca ggtcgttata gagatgcaag gcgtggtaag 4080
agagtttata gacagcagcc agccaccgct tagcgaggta aatcaggaga agtttaacaa 4140
tatcaagaaa gcactaagcg aagcgaaata actaaacctt catgcggcgg atttttccgc 4200
cgccattatt agcgagatag cgatgcacgt tacagccaag ccctccagtt ttcaatgtaa 4260
tctcaaatgt gattactggt ttaccttga aaaagagtcg cagtttactc atgaaaaatg 4320
gatggatgac agcactttga aagagttcat caaacaatat atcgacgcgt ctggcaatca 4380
gggtctatctt acctggcaag gcggtgaacc cactctggct ggccctggatt ttttccgtaa 4440
agttattcac tatcaacaac gctatgcagg ccaaaaacgt atttttaatg cattacaaac 4500
gaatggcatt ttattgaata atgaatgggtg tgccttctca aagaacatga atttctgggtg 4560
gtatctcgat cgatggcccc caggagttac atgaccgtta cagacgcagt aattcaggta 4620
acgggtacttt tgcaaaagtg atagcagcca tcgagcgtct gaaatcatat caagtagagt 4680
ttaatacgtt aaccgtcatt aataacgtta atgtocatta ccctcttgag gtttatcatt 4740
ttttaaaatc tatcggcagt aaacatatgc aatttatcga attgctagaa accgggacgc 4800
ogaatatiga tticagtggg catagtgaga acacattcgg tatcattgat ttttctgtgc 4860
ctcccacggc ttatggcaag tttatgtcaa ccatttttat gcaatgggtt aaaaacgatg 4920
tgggtgaaat ttcatccgt cagtttgaaa gctt 4954

```

<210> 13

<211> 3796

<212> DNA

<213> *Escherichia coli*

<220>

<223> Designated as EC-39

<400> 13

```

aagcttaatc gcgtgaatca ggagtaaaaa aatgacaacc cagactgtct ctggtcgccg 60

```

ttattttcaag aaagcgtggc tgatggagca gaaatcgctt atcgctctgc tgggtgctgat	120
cgcgattgtc tcgacgttaa gccogaactt tttcaccatc aataaacttat tcaatattct	180
ccagcaaacc tcagtgaacg ccattatggc ggtcgggatg acgctgggtga tcctgaacgc	240
gggcatcgac ttatcggtag gttctctgtt ggcgctgacc ggcgcagttg ctgcatctat	300
cgctggcatt gaagtcaatg cgctgggtggc tgtcgctgct gctctcgctg taggtgcgca	360
attgggtgcgg taaccgggggt gattgtagcg aaaggctcgcg tccaggcggtt tatcgctacg	420
ctgggttatga tgcttttact gcgcggcggtg accatgggtt ataccaacgg tagcccagtg	480
aataocgggt ttactgagaa cgccgatctg tttggctgggt ttgggtattgg tcgtccgctg	540
ggcgtaccga cgccagtcctg gatcatgggg attgtcttcc tcgcggcctg gtacatgctg	600
catcacacgc gtctggggcg ttacatctac gcgctggggc acaacgaagc gacaacgcgt	660
ctttctggta tcaacgtcaa taaaatcaaa atcatcgctt attctctttg tgggtctgctg	720
gcctcgctgg cgggatcata gaagtggcgc gtctctctcc cgcacaacca cggcggggac	780
tggctatgag ctggatgcta ttgctgcggt ggttctgggc ggtaacgagtc tggcggggcg	840
aaaaggctgc attgttggga cgttgatcgg cgcattaatt cttggcttcc ttaataatgg	900
attgaatttg ttaggtgttt cctctatta ccagatgac gtcaaagcgg tgggtgatttt	960
gctggcgggtg ctggtagaca acaaaaagca gtaataacga ctacaggcac atcttgaata	1020
tgaacatgaa aaaactggct accctgggtt ccgctgttgc gctaagcgc accgtcagtg	1080
cgaatgcgat ggcaaaagac accatgcgc tgggtggtct cagcttaac aaccgttct	1140
ttgtatcgct gaaagatggc gcgcagaaag aggcggataa acttggctat aacctgggtg	1200
tggactocca gaacaaccg gcgaaagagc tggogaacgt gcaggactta accgttcgcg	1260
gcacaaaaat totgctgatt aaccgaccg actccgacgc agtgggtaat gctgtgaaga	1320
tggctaacca ggogaacatc ccggttatca ctcttgaccg ccaggcaacg aaagggtgaag	1380
tggtgagcca cattgcttct gataacgtac tgggcggcaa aatcgctgggt gattacatcg	1440
cgaagaaagc ggggtgaagggt gccaaagtta tcgagctgca aggcattgct ggtacatccg	1500
cagcccgta acgtggcgaa ggcttcacgc aggcggttgc tgcacacaag tttaatgttc	1560
ttgccagcca gccagcagat tttgatcgca ttaaagggtt gaacgtaatg cagaacctgt	1620
tgaccgctca tcggatgtt caggctgtat tcgggcagaa tgatgaaatg gcgctgggcg	1680
cgctgcgcgc actgcaaact gcgggtaaat cggatgtgat ggtcgctgga tttgacggta	1740
caccggatgg cgaaaaagcg gtgaatgatg gcaaactagc agcgactatc gctcagctac	1800
ccgatcagat tggcgcgaaa ggcgtcgaaa ccgcagataa agtgctgaaa ggcgagaaa	1860
ttcaggctaa gtatccgggt gatctgaaac tgggtgttaa gcagtagttt taatcagggtt	1920
gtatgacctg atggtgacat aaatacgtca tcgacagatg aacgtgtaat ataaagaaaa	1980
gcagggaacg cgccacccta acacgggtggc gcattttatg gacatccga atatgcaaaa	2040
cgcaggcagc ctggtgttgc ttggcagcat taatgctgac cacattotta atcttcaatc	2100
ttttcctact ccaggcgaaa cgtaaccgggt aaccactatc aggttgcaat tggcggcaaa	2160
ggcgcgaatc aggtgtggc tgcgtggcggt agcgggtgca atatcgcggt tattgcctgt	2220
acgggtgatg acagcattgg tgagagcggt cgccagcagc tcgccactga taacattgat	2280

```

attactccgg tcagcgtgat caaaggcgaa tcaacagggtg tggcgtgat ttttgtaaat 2340
ggcgaagggtg agaatgtcat cgggtattcat gccggcgcta atgctgccct tccccggcg 2400
ctgggtggaag cgcaacgtga gcgtattgcc aacgcgtcag cattattaat gcagctggaa 2460
tcaccactcg aaagtgtgat ggcagcggcg aaaatcgccc atcaaaataa aaactatcgt 2520
tcgcttaacc cgctccggct cgcgaacttc ctgacgaact ctgcgctgtg gacattatta 2580
cgccaaacga aacggaagca gaaaagctca ccgggtattcg tgttgaaaat gatgaagatg 2640
cagcgaaggc ggcgcaggta cttcatgaaa aaggatatccg tactgtactg attactttag 2700
gaagtcgtgg tgtatgggct agcgtgaatg gtgaagggtca gcgcgttcct ggattccggg 2760
tgcaggctgt cgataccatt gctgccggag atacctttaa cggtgcgta atcaaggcat 2820
tgctggaaga aaaaccattg ccagaggcga ttctgtttgc ccatgctgcc gctgcgattg 2880
ccgtaacacg taaaggcgca caaccttccg taccgtggcg tgaagagatc gacgcatttt 2940
tagacaggca gaggtgacgc ttggctacaa tgaaagatgt tgcccgcctg gcggggcgttt 3000
ctacctcaac agttttctcac gttatcaata aagatcgctt cgtcagtga gcgattaccg 3060
caaagtgagc gcgattaaag actcaattac gcgccatcag ctctggcgcg tagcctcaaa 3120
ctcaatcaaa cacataccat tggcatgttg atcactgccca gtaccaatcc tttctattca 3180
gaactggtgc gtgtcgttga acgcagctgc ttcgaacgcg gttatagtct cgtcctttgc 3240
aataccgaag gcgatgaaca gcggatgaat cgcaatctgg aaacgctgat gcaaaaacgc 3300
gttgatggct tgctgttact gtgcaccgaa acgcatcaac cttcgcgtga aatcatgcaa 3360
cgttatccga cagtgcctac tgtgatgatg gactgggctc cgttcgatgg cgacagcgat 3420
cttattcagg ataactcgtt gctgggcgga gacttagcaa cgcaatatct gatcgataaa 3480
ggtcataccc gtatcgctg tattaccggc ccgctggata aaactccggc gcgctgcggt 3540
tggaagggtta tcgggcggcg atgaaacgtg cgggtctcaa cattcctgat ggctatgaag 3600
tactgggtga ttttgaattt aacggcgggt ttgacgctat gcgccaactg ctatcacatc 3660
cgctgcgtcc tcaggccgtc tttaccggaa atgacgctat ggctgttggc gtttaccagg 3720
cgttatatca ggcagagtta caggttccgc aggatatcgc ggtgattggc tatgacgata 3780
tcgaactggc aagctt 3796

```

<210> 14

<211> 6914

<212> DNA

<213> *Enterobacter cloacae*

<220>

<223> Designated as ET-49

<400> 14

```

aagctttttcg agttcgccat ccggcaacag ctactgagc ttttacgcgc ccagggtgcc 60

```

tttgaactca	attcccagct	cagtaaggcg	gtcctgaata	atctctttgc	gagatttttc	120
actggtaccg	gcatcaggtg	ttgcaggttt	cagctcgcca	ccagcctcgc	ccttcatcag	180
ccggacgtta	gacttcagcg	ccgggtgaag	atctttcaac	tccaccacgt	cgccaacctt	240
tacgccgaac	catggggcga	caacttcgta	tttagccatg	ctgtttcctt	acgccaggtt	300
agcgccgtag	acaacgccag	acaggcctga	tcgtctgcag	taatttgcag	gccttcagca	360
gacatgatct	ggaagttgta	gttaacgtta	ggcagtgggc	gcggcagtg	cacaacgcc	420
acagccatac	ccaccagtgg	ggagatcacg	tcacgacgac	gaacgtacgc	gataaactcg	480
ttaccgggtca	gcgcgaagtc	atgcggattt	ctttcaccgg	tgcgaaatggc	agaacagcct	540
gcaggagagt	gccgctcacc	acaccattaa	ctacgtatgg	ctgagccata	tttgcccaga	600
tctcagggga	aaccacatc	acatcatact	gagctacttt	gttggtgcgt	gcggtggtac	660
cgaatgctcc	tttaccaaag	aactcaaaat	attgagtcgt	ggttgcgctg	gtcaggtcga	720
tgttcgcacc	accagcacca	gaaccgaggt	taatcttctt	ggtgttgcg	tggttcttga	780
tgccctgcgc	cgggtaggac	tgaacctgaa	ttttgaatc	gccgttcagg	tagtagttga	840
cgcgcttctg	gttgaacttg	cgcattctcg	ccatctgcga	atccagaacc	agatcaatgc	900
ctacagagtt	aaggccagca	gcatgacgcc	agttaacacc	gtagccagca	gtgaacaccg	960
gaatcgggtc	gccatcgctc	gcgtagtcag	tgtggtcgaa	ggagaatggc	gcctgaccat	1020
cgatgcttac	tgacacgtcg	tcagcgatgt	cgccgaccac	gttatacagc	ttggcggttt	1080
taccaaccgg	cagcacggtc	tgaacgccga	tcaggtcgtt	tacgatttcc	atgccaaactt	1140
cctgatcccc	cagctgcagc	acctggttgt	caatctcagc	ccagaagtca	cgggagaaac	1200
cgccaacagc	gttacaagcc	agcatgtcag	gcgtcatcat	tgcgcggtta	gctgcaatga	1260
tggaatcggt	ctgtaggttc	cacatgttgc	ggtttgccca	cagctcactc	cagtgcctgc	1320
cgaggcggga	gttagtcgcc	agcgtctctt	tagagaagta	catatgtgtt	tgtccttttg	1380
ttacgcgcca	gctgcggcga	cagtgccaac	gcgcatacgc	acgcgaatga	agtcagtgg	1440
gctggccgcg	atggtgtatt	catcctggct	gtagccgatc	actgaatcag	tgtcggatgt	1500
ggcaagggtta	aactgaccgg	cagttcccag	cttgatcggg	ctgtcttttt	tatacgcacc	1560
aggcaggcag	cgcagcgcca	gctcacgacc	ttcttcgacg	tagttaccta	ctgccgaatc	1620
cccggcaggg	atttcttcgg	tgattgtcag	gccctggtga	taaccgacat	cgatgatgta	1680
caggcggccg	gttagcgcg	tggcctgagc	gaatttatcg	gatgagttga	tggttgcggc	1740
ggtgccagga	agcaaccgg	cggccgttgt	gcgggtttcg	gtcttgtaca	gagactgacc	1800
gtcgatatta	acgcgacgat	aacgtggcat	tattccggct	ccttacttga	agtgttcgtc	1860
tgcggtcgg	gcgcgggttt	ctttgtgctg	ctgagcattg	ttggtgccca	gcgacttgaa	1920
catcgcttcc	agagcttcgc	ctgacagagc	gttcgcgagc	gatatcgcca	tggaccttcg	1980
caaccgcttc	gcgctttgct	ttctcttcgg	cacgggagtt	cgcggtaagg	gtttccgcga	2040
gttgcttctg	attggcctgc	agcgcatcaa	ccttttcgcg	gagaggctta	atagccgctt	2100
cagtattgg	cgcaacagcc	tggccgatca	tgtgcgat	ttgttcag	tcttctttgg	2160
ttaaaggcat	gtgcctccg	ttttgtggtt	tggtgcaggc	tgttcctgcg	gtgtgaatag	2220
agctttgaat	tgtagcgac	gactgccacc	cacgaactct	ggcgcgctac	tgcggttccg	2280

gtatogtoga	ttgtgatott	cccgccatca	gcgaataaccg	taaacctgag	catcacccgc	2340
atttgcacg	atgaccacct	gcgagtcagt	gagtcagcaa	cccaggcata	ttcatccgtg	2400
cccgccgcaa	acttggtttt	ggctgcccga	tcgagacgct	gctcgcgctc	ccggtaggat	2460
tcacccacca	gcgcgccgga	gttcgcttta	agcggctgcg	ccagatcggc	gtttaccatc	2520
aggccaacgc	cctgctcagg	ggtggcggtt	ccgacttcgt	gcagtaggat	cgcgtogtgg	2580
tccatgctgt	gaatcttcgc	cacccactcg	gcacccgtag	ctctctgttg	ttcgttaggc	2640
tcaagctggt	cgaggaaagc	ggcgacactg	gtatgaatcg	gcggaacgtc	atcgccgcgc	2700
tcgatggctg	cgacgcgctc	aagtagttct	cggccacctt	cagactcacc	ggcgcgggca	2760
acatcaaccc	acttttgcag	gtagatacga	ttaccggact	tottaacgtt	gcggttccac	2820
gcgcgcgatat	ggcctgcgtt	aatccctcc	ggggagaaag	cagacacgaa	ctgaccatta	2880
acctgagggg	ggcccagcgg	cgcacgggta	ccttccagcc	ccttatagtg	ggcgctogatt	2940
tgtctttgcg	tgtacaagcc	gccattcatg	acgacgttag	ctggaagtgt	gtagctcggc	3000
agcaccagggt	gctcacgccc	gttgtatgtt	tcgcgcggga	tagactgggt	gttcaccttt	3060
gtggtgatgt	tgacctgaat	atgctcacca	tgtttcggtg	cctggattgg	acgctgtgct	3120
tcgtgggttta	cctggaattt	catgagttat	ttctccgccc	aggcgtaacc	gctcgccctgc	3180
atogatitatt	attcctgttt	gagtttcgtg	atggtgtccg	ggtattccgg	cttgccgtcc	3240
gcatccacca	gcaccgactg	ctggctgcac	ttgcagttga	tggagttgcc	atctttgctg	3300
taccagtcac	gcacctottc	gttggtgtag	aggtgggcat	gggcgcactg	cgtgggtatg	3360
tcgcgttgct	ggcgacagag	ctgagatgtg	aaccagcagc	gttttaaggc	cgaacaggtc	3420
attcgccctct	tgggtcttcat	cccacttggc	cgggcgcagc	gcggtagtca	cttcagtgcg	3480
tgtatccgg	ttagcccggc	gtttctcgat	gccggctctg	tctgtcaggt	tgcgggcaat	3540
gtccagagga	ttgagcccg	gcccaacacc	atcagtaaga	cacgcgccat	gtcgcgctta	3600
acgtcagccg	tcagcccctt	catttcctca	aatacacgcg	catgcaccag	cgccatgcgt	3660
ttctgatact	ggtcgcttgc	gaggatggag	gccagcgact	cacgcccggc	tgcgtacacc	3720
ggggattgct	gactgagggt	gtagaacgac	tgcccggtcc	ctttttccga	agccagatcg	3780
atgtactcgt	aaaaccacag	gtcgtaatcg	ccaccttcaa	gcagtacctg	atcaaccagg	3840
taactggcat	cgttcaggat	gatggagagt	agcattgggt	ttagctggta	ttcgatatctg	3900
gcgtttactg	cgaggggagga	aggatatttt	ttgagtgtcg	atttgtaacc	cttgccaatc	3960
ttattcatcc	gcctggcgaa	gttttccatt	gcccgccgtt	ccagcgcato	ggctccggtc	4020
ggatcctgat	agttacgcgg	cagaatcggt	ggcttcgtct	tttcgtcgc	catcctcttc	4080
tcctaattgga	aattcatoga	cgttttcata	accggcagca	gtgcggaatt	tcttcaagac	4140
taaaggctgg	ttttctccg	ctcccctgga	acgtctgggt	aatctctgcc	atgggttttg	4200
catttgcgag	tttctcagtt	ccagtctgtt	cgttgaggtc	atcccagata	accgtcttct	4260
cgtgactgc	atcaataatt	ttcagggtoga	tgagcttgct	actgaagtct	tcaatttoga	4320
atgacaggtc	accgcgcctg	gaatggcagc	gcgcgttgaa	atatttctga	tcctcgggtgc	4380
ttgccctttc	accgctctgc	atcccaacca	gaaccttcac	agggatataca	acagatgcag	4440
cgaagggttg	caggttgacg	ttatagggtcg	ctgacggatc	cgtacagct	gtgaccagtg	4500

gtgtgactgt	agcccccttg	gttgtcatca	gaacatcggt	accacgggtc	atttccccgg	4560
caacttcggt	aaacttatcc	tgcaactcgt	ccatgtcacg	ccataaagt	acgcgagatt	4620
gttgaaatcg	atttccttct	caaagttgac	attaagctgc	cgcgcggcgt	tctttaggaa	4680
tgactcacca	gaaccaccct	cgaccttctc	aaggctgacg	caggcggttat	agccaggctc	4740
aaggaagcca	atagcatcat	tagaatagtc	accaaggata	aagaocgat	cgggatgtac	4800
gaagcgtga	ttagttccac	cgcttggaag	gctctcaaca	tatttccact	gctttggctg	4860
cccgtagcct	gccgatttct	ggtcagttac	ccactcgctg	actgttaatg	accagccca	4920
tgcgatcgta	acctttttta	gtgacttgcc	acgaacaaca	ggctgatccc	atgttctgga	4980
atcattgata	tgacgcagga	tacccgcata	acgtccgacc	tgtcggcggc	ggtctgcttc	5040
agcaaaagcc	cgccaaaggc	gctttgtgaa	aacctttttg	gtgttcttct	cccaggcagt	5100
ttcatcctta	ctctcgtcgg	catcatcacc	ctcgatgatt	tccgggttgg	tctgccagca	5160
cttgcaccac	agcttctcta	ctgcgccgtg	ggctattcca	ccgcgacgat	acagtgcgta	5220
gaggttttcg	taagtgcact	gctcagggaa	tccatactcg	caccatgcgg	aatggcgctt	5280
attgtccagc	cccattgtag	gcgccaacag	ccccatacgg	gcacgggcca	tccgcgcctc	5340
gttcaacgca	tggttgacgg	cgagagttaa	tttgtcagtc	atggtttgtc	cgttggtgga	5400
tttaaggcat	aaaaaaaggc	cgctttggcg	accttgtggc	tatttaaaaa	gctaaaactct	5460
gttgaacgaa	ataaacataa	tctgctcagg	cttaacgcca	taatcacttg	ccaacttotg	5520
agtgcactca	attaagacag	ttgatgcaga	tttogaagag	cttgcaccat	aaatttcgaa	5580
gttttcaaat	actccgccgt	tggtgtggta	aatottatat	gacataaacc	aatcattcat	5640
aatatctact	cccttacaga	attgagtaga	tattatcggc	aagtgcataat	gtttctttta	5700
attatctcaa	ccttttcggg	atcatcatcc	cggccatctg	gcccttacgt	ttaatgtgtc	5760
cgtcgaggct	gtagcgaata	cgtcccagc	agtgttcgta	accgtctgcc	agtttaggca	5820
atacctcgcc	ggtgatgcgg	tccgttttgt	aggaccacat	gcgggcctct	ctcgccacat	5880
tcttgacgcg	aggatggata	atgatttcgt	caaagccgcg	aagatgcgcg	ataccgtcct	5940
caacactccc	ctgccatttc	tccgcagccg	agatgttgaa	gccctggcgc	ttgagatagc	6000
tgatagtctc	gggtcggggc	gagtgggcct	tgatgggcca	gtcacgcgat	ccggggattg	6060
tgctgtatag	ctctggcata	tggtcgagct	ctgtctgctg	accgtatgcc	tcttattcga	6120
tgtacagccg	gttgtgcagg	atgaacgagc	gcaccagcgt	gttagggctt	ttggcgaaac	6180
cgaagtcagc	acogaagaaa	aggcgatcgg	cctctttcca	tagctgggtc	gagaactcag	6240
cgatccggta	tttaccggcc	agcaactgct	tatcagagtt	ttcgaggtaa	gcaccttccc	6300
aaacccaogc	gtatgttgcc	gggtcaaggc	ggcgctgato	gttctgtcgc	tcaccttcca	6360
gcacgtcggg	gaaccatgga	ttatccgtgt	agttcatctc	aacgtgatac	agtogtcgcc	6420
agcctcttta	cggaaaogct	tatccgtgcg	ctgccgtcgc	gotccgggtt	ccatgtcacc	6480
caaactctct	aaccttctct	acgaacggtc	gggctcagct	tctgccaggc	tatttcgctg	6540
actgattcag	cctcatcaac	ccaacagagc	aagatgcgcg	ctttcgactt	gatgctgtcg	6600
aggttatgcc	gcagaccgca	gaacacgtag	ttaacgctct	tgctgatggg	goggatgtac	6660
ttctgcgcga	tatcaaagtt	ggaagccagc	cagggaaacag	acaggatagc	ctgtttccacc	6720

tcttgcatatc	tgcactcttc	cagtgagttc	atgaattcac	gcgcacagag	caccacgccg	6780
ctttcacggt	tcatcatcga	ctgatacgcc	tttacggctg	tcatcagcgc	aaaagtgcgc	6840
gtcttggcac	taccacgccc	accatgcgag	caccggtaac	gcttattctc	ggcgatgaac	6900
agtggcgcaa	gctt					6914

&lt;210&gt; 15

&lt;211&gt; 5975

&lt;212&gt; DNA

<213> *Klebsiella pneumoniae*

&lt;220&gt;

&lt;223&gt; Designated as KI-50

&lt;400&gt; 15

aagcttattc	cacgctggag	gcgtccggga	ttatcggcgt	caacgctatc	gccggcatcg	60
ccgggaccat	catcgccggc	atgctctccg	accgcttttt	caaacgcaac	cgcagcgtga	120
tggccggatt	catcagcctg	ctgaacaccg	ccggcttcgc	cctgatgctc	tggtcgccgc	180
acaattacta	cactgatatt	ctggcgatga	ttatcttcgg	ggccaccatt	ggcgctctga	240
cctgcttcct	tggcgggctg	atcgccgtcg	atatctcttc	gcgcaaggcc	gccggggccg	300
cgctcggcac	catcggcata	gcagctacgc	cggcgcgggc	ctgggcgagt	ttctcaccgg	360
gttcattatt	gataaaacgg	ctatccttga	aaacggcaaa	acgctgtatg	atttcagcac	420
gttggcgctg	ttctgggtgg	gtacggctctg	ggttcngcgc	tactctgttt	taccactgcc	480
gccatcgctg	cccggcgcca	tgccgtcgaa	cggcagacct	cgttctcctc	ataaccgatt	540
aacgaataag	gaagaagata	tgatgcctgc	aagacatcag	gggctgttac	gcctgtttat	600
cgctcgcgcg	ctgccgtgc	tggcgctgca	atctgccgcc	gccgcggact	ggcagctgga	660
gaaagtggtc	gagctcagcc	gccacggtat	togtccgccg	acggccggca	accgggaagc	720
catcgaggcc	gccaccggcc	gaccgtggac	cgagtggacc	acccatgacg	gggagctcac	780
cggccatggc	tatgccgcgc	tggtcaacaa	agggcgctgc	gaaggccagc	attaccgcca	840
gctcggcctg	ctgcaggccg	gatgcccgac	ggcggagctg	atatacgtgc	gcgccagccc	900
gctgcagcgg	acgcgagcga	ccgcccaggc	gctgggtggat	ggcgcccttc	ccggtgcggg	960
cgtegcctatc	cattatgtoa	gcggggatgc	cgatccctcg	tttcagaccg	acaagttcgc	1020
cgccacgcaa	accgaccccg	cccgccagct	ggcgcggtga	aagagaaggc	cggggatctg	1080
gcgcaggctg	gcaggcgctg	gcgcgcacca	tccagctatt	gaaacaggcg	gtttgtcagg	1140
ccgataagcc	ctgcccgcgc	ttcgataccc	cgtggcaggt	cgagcagagc	aaaagtggga	1200
agaccaccoat	tagoggactg	agcgtgatgg	ccaatatggg	ggagacgctg	cgtctcggct	1260
ggagtgaaaa	cctgcctctc	agccagctgg	cgtggggcaa	gatcaccocg	gccaggcgaga	1320
tcaccgcctt	gotgcgcgtg	ttaacggaaa	actacgatct	gagtaacgat	gtgttgtata	1380



ccgcgcacaaa	acgcgggtcg	gtgctgctca	acgctatgct	cgacggcgctc	aaaccggagc	1440
gaatcgaacg	tacgctggct	gctgctgggtg	gccatgacac	caatatcgcc	atgggtgcgca	1500
cgctgatgaa	ctttagctgg	cagctgccgg	gctacagccg	gggaaatata	ccgccggggca	1560
gcagcctggg	gctggagcgc	tggcgcaacg	cgaagagcgg	agaacgctat	ctgcgggtct	1620
atttcagggc	ccagggcctc	gacgacctgc	gtcgtctgca	gacgccggac	gcgcagaccc	1680
cgatgctgcg	tcaggagtgg	catcagccgg	gctgccgtca	gaccgatgtc	ggtagcctgt	1740
gtcccttcca	ggcggctatt	accgccctcg	gtcagcgctat	cgaccgatca	tccgccccgg	1800
cggtagcatg	gtcctgccgt	agcggcgcg	tgtttgtccg	ggccccggaa	aacctttttt	1860
tccaggccgg	cacgacgtcc	gttatccgtt	gtccggcgca	aacgccccgg	cgcgacacctg	1920
cgccgggggtg	acaccgctg	tccagcacc	agccgcttat	cagcccagca	ggcgtgacgt	1980
cgaacgcgg	attgtaaacg	gtggccccc	tggcgccca	ctgtaccgcg	ccgaagctgc	2040
ccgccactcc	ggtcacttcc	gccgccgcgc	gctgctcaat	ggggatcgcc	gccccgttcg	2100
ggcaatggcg	gtcgaggggtg	gtctgcgggg	cagcgacgta	aaacgggatc	tgggtgataat	2160
gggccaaaaac	cgccagagaa	taggtgccga	ttttattcgc	cacgtcgccg	ttggcggcga	2220
tacggtcggc	gccgaccac	accgcatcca	cctgccccctg	cgccatcagg	ctggcggccca	2280
ttgaatcggc	gatcagctga	tagggcacgc	ccagctcgcc	cagctcccag	gcggttaaac	2340
gaccgccctg	cagcagcggc	cggtttcat	caaccatac	gttggtcact	tttcctgccc	2400
ggtgcgccag	cgcgataacg	ccgagggcgg	tccctacccc	ggcggtcgcc	agggccaccgg	2460
tgttgacagt	ggtcagcagt	cgactgccgg	gcttcaccag	cgactgccc	gcctcagcga	2520
tgcggtcgca	cagctgttta	tcttcttcga	ccagacgcaa	ggcttcgcgt	tccagcgccct	2580
gcgggtaatc	tccgggccag	cgctgcttca	tgcgatcaga	ttattcatca	ggttgaccgc	2640
cgtcggccgc	gccgcgcgca	gtctccagcg	cctgctggag	tgcattcccg	ttcaggccgc	2700
gctgggccag	cagggccagc	agcaggctgg	cggacaggcc	aatcagcggc	gcgcgcgcga	2760
ccccgcagg	atgaatatgg	tccaccagca	gcgcaacgtt	atccgccgcc	agccagcgtt	2820
tttctgctgg	caaggcctgc	tggtcgagaa	taaaaagctg	attttcactc	accgcaggc	2880
tgggtgtctg	taatgtctgc	atgtcgttaa	atccctgttg	cgttggttga	tcacattgtg	2940
tcaggatgga	atccagaagt	atagacgtct	gaacggctta	atcagaattc	gaggatcgag	3000
gcaatgtcgc	aataccatac	cttcaccgcc	cacgatgccg	tggcttacgc	gcagagtttc	3060
gccggcatcg	acanccatct	gagctggtca	gcgcgcagga	agtgggcgat	ggcaactcaa	3120
tctggtgttt	aaagtgttcg	atcgccagg	cgtcacgggc	gatcgtcaaa	caggctctgc	3180
cctacgtgcg	ctgcgtcgcc	gaatcctggc	cgctgaccct	cgaccgcgcc	cgtctcgaag	3240
cgcagaccct	ggtcgcccac	tatcagcaca	gccgcgagca	cacggtaaaa	atccatcact	3300
ttgatcccg	gctggcggtg	atggtgatgg	aagatctttc	cgaccaccgc	atcttgccgc	3360
gagagcttat	cgctaacgtc	tactatcccc	aggcgccccg	ccagcttggc	gactatctgg	3420
cgcagggtgct	gtttcacacc	agcgatttct	acctccatcc	ccacgagaaa	aaggcgcagg	3480
tggcgaggtt	tattaacccg	gcgatgtgcg	agatcaccga	ggatctgttc	tttaacgacc	3540
cgtatcagat	ccacgagcgc	aataactacc	cggcgagcgt	gggaggccga	tgctcgccgc	3600

ctgcgcgacg	acgctcagct	taagctggcg	gtggcggcgc	tgaagcaccg	tttctttgcc	3660
catgcggaag	cgctgctgca	cggcgatata	cacagcgggt	cgatcttcgt	tgccgaaggc	3720
agcctgaagg	ccatcgacgc	cgagttcggc	tacttcggcc	ccattggctt	cgatatcggc	3780
accgccatcg	gcaacctgct	gcttaactac	tgcggcctgc	cgggccagct	cggcattcgc	3840
gatgccgccc	ccgcgcgcga	gcagcggctg	aacgacatcc	accagctgtg	gaccaccttt	3900
gccgagcgct	tccaggcgct	ggcggcggag	aaaacccgcg	acgcggcgct	ggcttacccc	3960
ggctatgcct	ccgcctttct	gaaaaagggtg	tgggcggacg	cggtcggctt	ctgcggcagc	4020
gaactgatcc	gccgcagcgt	cggactgtcg	cacgtcgcgg	atatcgacac	tatccaggac	4080
gacgccatgc	gtcatgagtg	cctgcgccac	gccattaccc	tgggcagagc	gctgatcgtg	4140
ctggccgagc	gtatcgacag	cgtcgacgag	ctgctggcgn	gggtacgcca	gtacagctga	4200
gtgcgcctgt	ttccctcacc	ccaacctctt	cccacaggga	gagggagcac	cccctaaaaa	4260
agtgccattt	tctgggattg	cccggcgngn	tgcgcttgcc	gggcctacag	atagccgcat	4320
aacggtttga	tcttgcactc	tttcgtaggc	cgggtaaggc	gaaagccgcc	acccggcaga	4380
catgcgagta	caattttgca	tttaccttac	cctcacccca	gatactcaat	caccgatagc	4440
ccgccgttgt	aatcggtgct	gtagataatg	ccttgcgcgt	cgacaaacac	gtcacaggac	4500
tggatcacc	gcgggcggcc	gggacgggta	tccatcattc	tctcagcgca	gccggcacca	4560
gcgccccggt	ctccagcggg	cgatacgggt	tggaaatgtc	gtaagcccg	acgccggcat	4620
tctgatacgt	ggcaaaaatc	agcgttgagc	tgacaaagct	ccccggccgg	ttctcatgca	4680
ggttgtgcgg	accgaaatgc	gcccctttcg	ccacgtaatc	cgcttcatcc	ggcggcggga	4740
aggtggcgat	gctcaccggg	ttggttggct	cgcggatata	aaacagccag	atcagcttct	4800
cgcggtcctc	ctggttatcg	agcaccgctt	catccagcac	caccagcaga	tgcgatccg	4860
gcagcggcag	cgcggtatgc	gttcgcgcgc	cgaacggcgg	gctccagttg	cgatggctaa	4920
tcagcctcgg	ctgggtacgg	tctttgacat	ccagcagcgt	caggccggcg	tgcgcccagc	4980
tgcgtaggcg	tatccccggc	aataatggcg	tgatgcagcg	catagcgttt	gccttgcggc	5040
cagtcgggtg	tttcaccgcc	cgcttgggtg	atccccggca	gccaccagcg	cccggtact	5100
tcgggcttac	gcggatcggc	cagatcgatg	gtcaggaaga	tgtagtcggt	aaaaccgtcg	5160
atcagcgcag	acacatacgc	ccagcgcgcc	ccgacgtacc	agatgcgggtg	aataccgatg	5220
ccgttaagcg	acaggaaact	gatttcccgc	gctgcgcggg	agtggaaata	tcaaagatgc	5280
gcagcccggc	gctccagccc	ctgtcctgca	catcgctgac	cgtgtcacc	accgagcggg	5340
tgtagtacac	cttctcatca	gcaaaacggg	cgtcagcaaa	cagatcccgg	gcgttgatca	5400
ccagcagcag	atcgtcatgc	gcctggagtg	cacgttccag	gtgcccggcg	gcgcggcaat	5460
atagttgacg	gtggtgggccc	gggtcggatc	gcgaacatcg	accacggaaa	aaccctgcga	5520
caccatatgg	ccgatatagg	cgaatccgcg	gtgcaccatc	agctgcacgc	cgtccggacg	5580
accgccctga	tcgctatggc	caatcagccg	catattgcgg	ctgtattcgg	gggaaggtaa	5640
tgetgacata	ggggatccct	ctcgcccgggt	ggcatggttt	ttccccctct	cctgcggaga	5700
gggcccgggg	gagggcacca	ggccgcgcgc	caccgccacc	cggcttgatt	ttatttgttc	5760
ttcgcttcca	gcgtcgcgaa	ccacggcgcg	ataaagtctt	cggctctggc	ccagccaggg	5820

ataattttcc ccagcgacgc cacgtttacc gctcccggct gggccgccag cagcgccctgg 5880  
 ggaatcgctg ccgccttgaa gtcgtaggtg gctggcgctg gctcgccggc gatcttggtg 5940  
 gcgatcagcc gcacgttggt cgcgccgata agctt 5975

<210> 16

<211> 899

<212> DNA

<213> *Candida albicans*

<220>

<223> Designated as CA-26

<400> 16

gaattcctag taagcgcaag tcatacagctt gcgttgatta cgtccctgcc ctttgtacac 60  
 accgcccgtc gctactaccg attgaatggc ttagtgaggc ctccggattg gtttaggaaa 120  
 gggggcaacc tcattctgga accgagaagc tgggtcaaact tggtcattta gaggaagtaa 180  
 aagtcgtaac aaggtttccg tagtgaacct gcggaaggat cattactgat ttgcttaatt 240  
 gcaccacatg tgtttttctt tgaacaaaact tgctttgcgg tgggcccagc ctgccgccag 300  
 aggtctaaac ttacaaccaa ttttttatca acttgtcaca ccagattatt acttaatagt 360  
 caaacttcaa caaacggatc tcttggttct cgcagcgaaa tgcgatacgt aatatgaatt 420  
 gcagatattc gtgaatcacc gaatctttga acgcacattg cgcctctctg tattccggag 480  
 ggcatgcctg tttgagcgtc gtttctccct caaacccgtg ggtttggtgt tgagcaatac 540  
 gacttgggtt tgcttgaaag acggtagtgg taaggcggga tcgtttgaca atggcttagg 600  
 tctaaccaaa aacattgctt gcggcggtaa cgtccaccac gtatatcttc aaactttgac 660  
 ctcaaatacag gtaggactac ccgctgaact taagcatatc aataagcgga ggaaaagaaa 720  
 ccaacaggga ttgcctcagt agcggcgagt gaagcggcaa aagctcaaatt ttgaaatctg 780  
 gcgtcttttg cgtccgagtt gtaatttgaa gaaggtatct ttgggcccgg ctcttgtcta 840  
 tgttcccttg aacaggacgt cacagagggt gagaatcccg tgcgatgaga tgaccggg 899

<210> 17

<211> 189

<212> DNA

<213> *Candida albicans*

<220>

<223> Designated as CA-26-1

&lt;400&gt; 17

gcgcaagtca tcagcttgcg ttgattacgt ccctgccctt tgtacacacc gcccgtcgct	60
actaccgatt gaatggctta gtgaggcctc cggattgggt taggaaaggg ggcaacctca	120
ttctggaacc gagaagctgg tcaaacttgg tcatttagag gaagtaaaag tcgtaacaag	180
gtttccgta	189

&lt;210&gt; 18

&lt;211&gt; 224

&lt;212&gt; DNA

&lt;213&gt; Candida albicans

&lt;220&gt;

&lt;223&gt; Designated as CA-26-2

&lt;400&gt; 18

gctgggtttg gtgttgagca atacgaacttg ggtttgcttg aaagacggta gtggtaaggc	60
gggatcgttt gacaatggct taggtctaac caaaaacatt gcttgcgggcg gtaacgtcca	120
ccacgtatat cttcaaactt tgacctcaaa tcaggtagga ctaccgctg aacttaagca	180
tatcaataag cggaggaaaa gaaaccaaca gggattgcct cagt	224

&lt;210&gt; 19

&lt;211&gt; 369

&lt;212&gt; DNA

&lt;213&gt; Candida albicans

&lt;220&gt;

&lt;223&gt; Designated as CA-26-3

&lt;400&gt; 19

aaacggatct cttggttctc gcagcgaaat gcgatacgta atatgaattg cagatattcg	60
tgaatcatcg aatctttgaa cgcacattgc gccctctggt attccggagg gcatgcctgt	120
ttgagcgctcg tttctccctc aaaccgctgg gtttggtgtt gagcaatacg acttgggttt	180
gcttgaaaga cggtagtggt aaggcgggat cgtttgacaa tggcttaggt ctaacaaaa	240
acattgcttg cggcggtaac gtccaccacg tatactttca aactttgacc tcaaatcagg	300
taggactacc cgctgaactt aagcatatca ataagcggag gaaaagaaac caacagggat	360
tgccctcagt	369

Application number / numéro de demande: JP02/05107

Figures: \_\_\_\_\_

Pages: 1/2/5/6/7/10/11

Unscannable items  
received with this application  
(Request original documents in File Prep. Section on the 10<sup>th</sup> floor)

Documents reçu avec cette demande ne pouvant être balayés  
(Commander les documents originaux dans la section de préparation des dossiers au  
10<sup>ème</sup> étage)

3/11

FIGURE 3






FIGURE 4



8/11

FIGURE 8



Subjected Digested Samples

SA : SA Digested Sample

SE : SE Digested Sample

PA : PA Digested Sample


EF : EF Digested Sample

EK : EK Digested Sample



9/11

FIGURE 9



SA : SA Digested Sample

SE : SE Digested Sample

PA : PA Digested Sample

EF : EF Digested Sample

EK : EK Digested Sample